

| Almonds, Pistachios & Walnuts |

## Senstar® Insecticide: Boost Your Insect Control

Senstar® Insecticide brings two effective modes of action for long-lasting control of soft-bodied insects, including mealybugs in pistachios, scale in walnuts and mealybugs and scale in almonds. With contact, translaminar and systemic activity that affects all life stages of targeted pests for long-lasting control, *Senstar* helps ensure high-quality tree nuts from the start.


- ▶ Controls target pests at all life stages, including eggs and crawlers
- ▶ Provides dual systemic activity to control pests moving into new vegetative growth as well as pests which populate in already developed foliage
- ▶ Delivers translaminar movement to reach target pests that feed on the underside of leaves
- ▶ Selectively targets harmful insect pests with minimal impact on beneficial arthropods

## Senstar is a Superior Management Tool for Gill’s Mealybug in Pistachios

*Senstar* effectively controlled the occurrence of Gill’s mealybug, as well as the severity of the population, better than Movento®.

Means followed by same letter or symbol do not significantly differ (P=0.05, LSD).

Pistachio variety: Kerman  
Location: Wasco, CA  
Source: Sawtooth Ag Research



## How To Use

<b>Rate</b>	12–18 fl oz/A
<b>Timing</b>	Almonds: After petal fall–best after full canopy leaf out Pistachios: After nut set at first sign of Gill’s mealybug crawlers Walnuts: After nut set
<b>Method</b>	Ground application. Only whole fields should be treated.
<b>Spray Volume</b>	100–400 gallons/A. Coverage is essential for good control. Apply in sufficient water volumes according to the tree size. Use of higher water volumes will assure better coverage and may provide better insect control.
<b>Adjuvant</b>	Use a penetrator type adjuvant. Do not use sticker/spreader type adjuvants.
<b>REI / PHI</b>	24 hours / 21 days

## Other Important Information

- ▶ Do not apply more than 18 fl oz/A per application
- ▶ Do not make more than two applications per year
- ▶ Do not apply more than 36 fl oz/A per year
- ▶ Minimum interval between applications is 14 days
- ▶ For control of San Jose scale, apply immediately after petal fall; under heavy infestation pressure or where difficult control conditions exist, a second application may be necessary
- ▶ For pistachios, the U.S. tolerance for pyriproxyfen is 0.02 ppm and spirotetramat is 0.25 ppm. To get the latest global MRLs, please refer to <https://bcglobal.bryantchristie.com/db#/pesticides/query> for both pyriproxyfen and spirotetramat tolerances by crop and export country.

