



NipsIt INSIDE[®] Insecticide

TREATED SEED

This seed has been treated with NipsIt INSIDE[®] Insecticide which contains clothianidin at the rate recommended by the manufacturer.

CAUTION **DO NOT USE TREATED SEEDS** **FOR FEED, FOOD OR OIL PROCESSING**

STORE AWAY FROM FEEDS AND
OTHER FOODSTUFFS
KEEP OUT OF REACH OF CHILDREN

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear long-sleeved shirt, long pants and waterproof gloves when handling treated seed.

FIRST AID

IF SWALLOWED:

Call a poison control center or doctor immediately for treatment advice.

Have a person sip a glass of water if able to swallow.

Do not induce vomiting unless told to do so by a poison control center or doctor.

Do not give anything to an unconscious person.

Have product container or label with you when calling a poison control center or doctor, or going for treatment.

PLANT BACK RESTRICTIONS

- Areas planted with treated seed of cereal grains (except rice, sorghum, sweet corn, wild rice, field corn and popcorn), sugar beet and soybean may be replanted immediately with bulb vegetables, canola, cereal grains (except rice), cotton, corn (field, pop and sweet), fruiting vegetables (including cucurbits), leafy vegetables (including brassica vegetables), rapeseed, root and tuber vegetables, sorghum, soybean and sugar beet. These areas may also be replanted after 30 days with dry beans, grasses, non-grass animal feeds and rice. These areas may also be replanted after 8 months with sugarcane. Do not plant any other crop in the treated area for at least one year after these treated seeds are planted.
- Areas planted with treated seed of canola, rapeseed and sorghum may be replanted immediately with bulb vegetables, canola, cereal grains (except rice), cotton, corn (field, pop and sweet), fruiting vegetables (including cucurbits), leafy vegetables (including brassica vegetables), rapeseed, root and tuber vegetables, sorghum, soybean and sugar beet. These areas may also be replanted after 30 days with dry beans, grasses, non-grass animal feeds and rice. Do not plant any other crops in the treated area for at least 4 months after these treated seeds are planted.

NOTE:

- Rapeseed greens and seed grown or harvested from *NipsIt INSIDE*-treated seed must not be used for feed or human consumption. Rapeseed grown and harvested from *NipsIt INSIDE*-treated seed is only for industrial uses and cannot be used for edible oil or any other human/feed consumption.
- Do not graze or feed soybean forage and hay to livestock.
- Cover or incorporate spilled treated seeds.
- Dispose of all excess treated seed. Left over treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements. Do not contaminate water bodies when disposing of planting equipment washwaters.
- Dispose of seed packaging in accordance with local requirements.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading.
- Regardless of the type of application (seed treatment, soil or foliar) do not apply more than a total of 6 oz clothianidin (active ingredient) per acre per cropping cycle except for soybean. Regardless of the type of application (seed treatment, soil or foliar) do not apply more than a total of 3.2 oz clothianidin (active ingredient) per acre per cropping cycle for soybean.
- Treated seed does not flow as freely as untreated seed. Make adjustments to compensate for this restricted seed flow to prevent lower than desired seeding rates.



The source of clothianidin, *NipsIt INSIDE*, EPA Reg. 59639-151

NipsIt INSIDE is a registered trademark of Valent U.S.A. Corporation

For **PRODUCT USE** information:
Call 1-800-6-VALENT (1-800-682-5368)
IN CASE OF EMERGENCY ONLY
Call 24 Hours A Day 1-800-892-0099

Valent U.S.A. Corporation
P.O. Box 8025
Walnut Creek, California 94596-8025

Form 1719-B