

Supplemental Label



Valent U.S.A. Corporation
1333 N. California Blvd., Ste. 600
Walnut Creek, CA 94596-8025

RESOURCE[®] HERBICIDE

FIELD CORN AND SOYBEANS

EPA Reg. No. 59639-82

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL AND PAMPHLET. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

DIRECTIONS FOR AERIAL APPLICATION AS PART OF A BURNDOWN PROGRAM AND FOR POSTEMERGENCE USE IN FIELD CORN AND SOYBEAN

RECOMMENDATIONS FOR AERIAL APPLICATION

GENERAL RESTRICTIONS

- Do not apply during low-level inversion conditions, when winds are gusty or under any other condition that favors drift.
- Do not spray when drift is possible or when wind velocity is more than 10 mph.
- Drift may cause damage to any vegetation contacted to which application is not intended. To prevent injury to adjacent desirable vegetation, appropriate buffer zones must be maintained.
- Do not apply this product within 40 feet of non-target plants including non-target crops.
- Do not apply this product within 70 feet of streams, wetlands, marshes, ponds, lakes, and reservoirs.

GENERAL INFORMATION

To obtain satisfactory weed control with aerial application of RESOURCE, use as part of a labeled tank mix. Uniform coverage must be obtained. To obtain satisfactory application and avoid drift, the following directions must be observed:

Maximum Use Rate

The maximum rate of RESOURCE that can be applied by air is 4 fl. oz./A. Do not apply more than an additional 4 fl. oz./A. of RESOURCE following an aerial application to corn or 8 fl. oz./A of RESOURCE following an aerial application to soybeans.

When applied aerially followed by ground application, do not exceed a maximum of 8 fl. oz./A. of RESOURCE to corn or 12 fl. oz./A of RESOURCE to soybeans per season.

Carrier Volume and Spray Pressure

Use RESOURCE in 7 to 10 gals. of water per acre and a maximum spray pressure of 40 PSI. Application at less than 7 gals. per acre will provide inadequate control. The higher gallonage applications generally afford more consistent weed control.

Nozzle and Nozzle Orientation

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray as fine droplets. Use the largest droplet size possible that provides sufficient coverage and control. Use nozzles which produce flat or hollow cone spray patterns. Use non-drip type nozzles, such as diaphragm-type nozzles to avoid unwanted discharge of spray solution.

Do not angle nozzles forward into the air stream and do not increase spray volume by increasing nozzle pressure. The nozzles must be directed toward the rear of the aircraft, at an angle between 0 and 15° downward. Do not place nozzles on the outer 25% of the wings or rotors.

Adjuvants and Drift Control Additives

Refer to tank mix partners label for adjuvant recommendation. Drift control additives may be used. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION. PLEASE REFER TO CONTAINER LABEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

Copyright© 2002 by Valent U.S.A. Corporation

Manufactured for
Valent U.S.A. Corporation
Walnut Creek, CA 94596-8025
www.valent.com
Form 2002-RES-0010
Made in U.S.A.

Products That Work, From People Who Care™

2002-RES-0010 2/02

Valent U.S.A. Corporation
1-800-6-VALENT (682-5368)
www.valent.com

