

# Supplemental Label



EPA Reg. No. 59639-99  
(EXCEPT NEW YORK)

## CHATEAU® HERBICIDE SW USE ON: STRAWBERRY

### Shielded and Hooded Application in Strawberry Row Middles

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.**

#### GENERAL INFORMATION

*Chateau SW*, at 3 oz per acre, can be applied to the soil a minimum of 30 days prior to transplanting strawberries provided the strawberries will be transplanted through a plastic mulch.

*Chateau SW*, at 3 oz per acre, can be applied with a shielded or hooded sprayer for the preemergence control of the weeds listed in Table 1, Broadleaf Weeds Controlled by Residual Activity of *Chateau SW*, in strawberry row middles.

Application Method	Minimum Time From Application to Harvest (PHI)	Use Rate Per Acre Per Application (oz)	Use Rate Per Acre Per Year (oz)	Special Use Instructions
Pre-transplant	Not applicable	3	3	Apply a minimum of 30 days prior to transplanting and prior to plastic mulch being laid. Apply as part of a tank mix to control emerged weeds.
Hooded or shielded sprayer application to row middles	Do not apply after fruit set	3	3	<b>Apply only to row middles – do not apply over strawberries.</b> Apply prior to weed emergence. Crop spotting may occur if an adjuvant is added. <b>Application after fruit set may result in spotting of fruit and should be avoided.</b> Do not allow spray drift to come in contact with fruit or foliage

#### SPRAY DRIFT MANAGEMENT

Avoiding spray drift at the application site is the responsibility of the applicator and the grower. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and grower

must consider the interaction of equipment and weather related factors to ensure that the potential for drift to sensitive non-target plants is minimal.

**Table 1. Broadleaf Weeds Controlled by Residual Activity of Chateau SW**

Common Name	Scientific Name	Organic Matter	Soil Type	Chateau SW Rates
Coffee Senna Common Ragweed Florida Beggarweed Golden Crownbeard Hairy Indigo	<i>Cassia occidentalis</i> <i>Ambrosia artemisiifolia</i> <i>Desmodium tortuosum</i> <i>Verbesina encelioides</i> <i>Indigofera hirsuta</i>	Up to 3%	All Soil Types	3 oz/A
Hemp Sesbania Jimsonweed Kochia Morningglories <sup>1</sup> Entireleaf  Ivyleaf Red/Scarlet Tall Mustard, Wild Palmer Amaranth Spurred Anoda Tropic Croton Waterhemp Common Tall Wild Poinsettia	<i>Sesbania exaltata</i> <i>Datura stramonium</i> <i>Kochia scoparia</i>  <i>Ipomoea hederacea</i> var. <i>integriuscula</i>  <i>Ipomoea hederacea</i> <i>Ipomoea coccinea</i> <i>Ipomoea purpurea</i> <i>Brassica kaber</i> <i>Amaranthus palmeri</i>	3 to 5%	Coarse and Medium Soils: (sandy loam, loamy sand, loamy silt-loam, silt, sandy clay, sandy clay loam)	3 oz/A
Carpetweed Chickweeds Common Mouseear Dandelion Eclipta Eveningprimrose, Cutleaf Florida Pusley Henbit Lambsquarters, Common Little Mallow Marestail/Horseweed Nightshades Black Eastern Black Hairy Pigweeds Redroot Smooth Spiny Amaranth Tumble Prickly Sida (Teaweed) Puncturevine Purslane, Common Radish, Wild Redmaids Shepherd's-Purse Smallflower Morningglory Spotted Spurge Venice Mallow	<i>Anoda cristata</i> <i>Croton glandulosus</i>  <i>Amaranthus rudis</i> <i>Amaranthus tuberculatus</i> <i>Euphorbia heterophylla</i> <i>Mollugo verticillata</i>  <i>Stellaria media</i> <i>Cerastium vulgatum</i> <i>Taraxacum officinale</i> <i>Eclipta prostrata</i> <i>Oenothera laciniata</i> <i>Richardia scabra</i> <i>Lamium amplexicaule</i> <i>Chenopodium album</i> <i>Malva parviflora</i> <i>Conyza canadensis</i>  <i>Solanum nigrum</i> <i>Solanum ptycanthum</i> <i>Solanum sarrachoides</i>  <i>Amaranthus retroflexus</i> <i>Amaranthus hybridus</i> <i>Amaranthus spinosus</i> <i>Amaranthus albus</i> <i>Sida spinosa</i> <i>Tribulus terrestris</i> <i>Portulaca oleracea</i> <i>Raphanus raphanistrum</i> <i>Calandrinia ciliata</i> var. <i>menziessii</i> <i>Capsella bursa-pastoris</i> <i>Jacquemontia tamnifolia</i> <i>Euphorbia maculata</i> <i>Hibiscus trionum</i>	Up to 5%	Fine Soils: (silty clay, silty clay loam, clay, clay loam)	3 oz/A

<sup>1</sup>Morningglory species are not adequately controlled on fine soils or soils with greater than 3% organic matter.

**THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. PLEASE REFER TO CONTAINER LABEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS. FOLLOW ALL APPLICATION DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED LABEL.**

**PLEASE CONTACT VALENT U.S.A. CORPORATION AT 800-6-VALENT (682-5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE.**

Manufactured for  
Valent U.S.A. Corporation  
P. O. Box 8025  
Walnut Creek, CA 94596-8025  
Made in U.S.A.



*Products That Work, From People Who Care*® | [www.valent.com](http://www.valent.com) | 800-6-VALENT (682-5368)

***Read and follow the label instructions before using.***

*Chateau and Products That Work, From People Who Care* are registered trademarks of Valent U.S.A. Corporation. © 2007 Valent U.S.A. Corporation. All rights reserved. Printed in the USA. 2007-CHSW-0010 XNY 1/08 mf/AV