

Supplemental Label



EPA Reg. No. 59639-119

CHATEAU® HERBICIDE WDG: FOR USE ON POME FRUIT, STONE FRUIT AND NON-BEARING FRUIT TREES REVISED USE PRECAUTIONS

**(For Use in All States Where This Product is Registered, Except Idaho and New York)
(Limited Use in Oregon and Washington)**

For Oregon and Washington, Use Only in the Following Counties West of the Cascade Mountains:
Oregon: Benton, Clackamas, Clatsop, Columbia, Coos, Curry, Douglas, Jackson, Josephine, Lane, Lincoln, Linn, Marion, Multnomah, Polk, Tillamook, Yamhill and Washington.
Washington: Clallam, Cowlitz, Grays Harbor, King, Jefferson, Kitsap, Lewis, Pacific, Pierce, Skagit, Snohomish, Thurston, Wahkiakum, Whatcom.

THIS SUPPLEMENTAL LABEL SUPERCEDES PREVIOUS LABELING FOR POME FRUIT, STONE FRUIT AND NON-BEARING FRUIT TREES.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

GENERAL RESTRICTIONS AND LIMITATIONS

- Do not apply more than 12 oz of *Chateau* WDG per acre during a single application.
- Do not apply more than 24 oz of *Chateau* WDG per acre during a 12 month period.
- Do not make a sequential application within 30 days of the first application.
- A maximum *Chateau* WDG rate of 6 oz/A per application should be used on any soil that has a sand plus gravel content over 80% if trees are less than 3 years of age.
- Do not apply to farm alleys or roads where traffic may result in treated dust settling onto crops or other desirable vegetation.
- Do not apply within 300 yards of non-dormant pears.
- Do not apply to powdery soils or soils that are susceptible to wind displacement unless irrigation can be applied immediately after application.
- Do not mow treated areas between bud break and final harvest. Dust created by mowing may drift onto desirable vegetation resulting in injury.
- Follow the most restrictive label limitations and precautions of the tank mix product(s) being used.
- Avoid direct or indirect spray contact to foliage and green bark (non-barked trunk with the exception of undesirable suckers).

For non-bearing fruit trees, *Chateau* WDG should be applied as a uniform band directed at the base of the trunk. For pome fruit and stone fruit, *Chateau* WDG can only be applied as a uniform band directed at the base of the trunk prior to “pink bud” in apple and “bud break” in stone fruit and pear. The preferred application timing for *Chateau* WDG is in the fall to maximize the potential for rainfall to activate and set the herbicide. Do not apply over the top of crop or allow spray to come in contact with crop as a result of application or drift.

Preemergence Application

Apply 6 to 12 oz (0.188 to 0.38 lb ai/A) of *Chateau* WDG per broadcast acre as a preemergence application. Preemergence (to weed emergence) applications of *Chateau* WDG should be made to a weed-free soil surface. Preemergence applications of *Chateau* WDG must be completed prior to weed emergence. Moisture is necessary to activate *Chateau* WDG on soil for residual weed control. Dry weather following application of *Chateau* WDG may reduce effectiveness. However, when adequate moisture is received after dry conditions, *Chateau* WDG will control susceptible germinating weeds.

Postemergence Application

Apply 6 to 12 oz (0.188 to 0.38 lb ai/A) of *Chateau* WDG per broadcast acre plus an adjuvant (0.25% v/v non-ionic surfactant or 1 qt/A crop oil concentrate). The addition of an adjuvant enhances *Chateau* WDG activity on emerged weeds. Thorough spray coverage is necessary to maximize the postemergence activity of *Chateau* WDG.

Refer to Table 1 for weeds controlled by the residual activity of *Chateau* WDG. *Chateau* WDG should be tank mixed with a labeled burndown herbicide for control of the emerged weeds listed in Table 2. Refer to tank mix partner’s label for additional weed species and increased weed heights claimed. Refer to tank mix partner’s label for additional restrictions, including minimum carrier volume and crops in which tank mix partner may be used. Burndown tank mix partners include glyphosate, paraquat, 2,4-D and Rely®. Tank mixes with glyphosate or 2,4-D containing products are not recommended during the period after bloom through final harvest to ensure crop safety from drift.

Residual weed control will be reduced if vegetation prevents the *Chateau* WDG from reaching the soil surface. If vegetation is heavy, it is recommended to use a burndown herbicide with *Chateau* WDG and make a sequential *Chateau* WDG application prior to the emergence of new weeds.

Carrier Volume and Spray Pressure

To ensure thorough coverage in burndown applications, use a minimum of 15 gallons of spray solution per acre. Use higher gallonage if dense vegetation or heavy crop residue is present.

Nozzle selection should meet manufacturer's gallonage and pressure recommendations.

Banded Application

Rates listed in Table 2 refer to a broadcast application covering the entire acre. When making a banded application, the rate must be reduced according to the following formula:

| | | | | |
|---|---|--|---|----------------------------|
| Amount Needed per Acre for Banded Application | = | $\frac{\text{Band Width in inches}}{\text{Row Width in inches}}$ | X | Rate per Broadcast Acre |
|---|---|--|---|----------------------------|

USE PRECAUTIONS FOR POME FRUIT AND STONE FRUIT

Pome Fruit: Apple, Crabapple, Loquat, Mayhaw, Pear, Pear (oriental) and Quince

Stone Fruit: Apricot, Cherries (Sweet and Tart), Nectarine, Peach, Plum (Chickasaw, Damson, Japanese), Plumcot and Prune

- For stone fruit and pear, apply only between final harvest and bud break.
- For apple, apply only between final harvest and pink bud.
- For pome fruit and stone fruit do not apply to row middles (area between berms)
- Do not apply within 60 days of harvest.
- Do not apply to trees established less than one year, unless protected from spray contact by non-porous wraps, grow tubes or waxed containers.

USE PRECAUTIONS FOR NON-BEARING FRUIT TREES

Non-Bearing Avocado, Fig, Grapefruit, Lemon, Olive, Orange, Pomegranate and Tangerine

- Do not apply more than 12 oz of *Chateau* WDG per acre during a single application.
- Do not apply more than 24 oz of *Chateau* WDG per acre during a 12 month period.
- Do not harvest fruit from treated trees within one year of application.
- Do not apply to trees established less than one year, unless protected from spray contact by non-porous wraps, grow tubes or waxed containers.
- Do not apply during the period after flowering through leaf drop, unless using shielded application equipment and the applicator can ensure spray drift will not come in contact with the crop foliage.

ADDITIONAL RESIDUAL WEED CONTROL

Chateau WDG may be tank mixed with oryzalin (Surflan[®]), simazine or diuron for additional residual weed control. Always read and follow label use directions for all products being used.

Table 1. Weeds Controlled by Preemergence Application of *Chateau* WDG

| BROADLEAF WEED SPECIES | | | | |
|-------------------------------|---|------------------------|-----------------------------|---------------------------|
| COMMON NAME | SCIENTIFIC NAME | ORGANIC MATTER | SOIL TYPE | CHATEAU WDG RATE |
| Bristly Starbur | <i>Acanthospermum hispidum</i> | Up to 10% ¹ | All Soil Types ² | 6 to 12 oz/A ² |
| Carpetweed | <i>Mollugo verticillata</i> | | | |
| Chickweeds | | | | |
| Common | <i>Stellaria media</i> | | | |
| Mouseear | <i>Cerastium vulgatum</i> | | | |
| Coffee Senna | <i>Cassia occidentalis</i> | | | |
| Dandelion | <i>Taraxacum officinale</i> | | | |
| Eclipta | <i>Eclipta prostrata</i> | | | |
| Eveningprimrose, Cutleaf | <i>Oenothera laciniata</i> | | | |
| Filaree | | | | |
| Redstem | <i>Erodium cicutarium</i> | | | |
| Whitestem | <i>Erodium moschatum</i> | | | |
| Fleabane | <i>Erigeron</i> spp. | | | |
| Florida Beggarweed | <i>Desmodium tortuosum</i> | | | |
| Florida Pusley | <i>Richardia scabra</i> | | | |
| Golden Crownbeard | <i>Verbesina encelioides</i> | | | |
| Hairy Indigo | <i>Indigofera hirsute</i> | | | |
| Hemp Sesbania | <i>Sesbania exaltata</i> | | | |
| Henbit | <i>Lamium amplexicaule</i> | | | |
| Jimsonweed | <i>Datura stramonium</i> | | | |
| Kochia | <i>Kochia scoparia</i> | | | |
| Lambsquarters, Common | <i>Chenopodium album</i> | | | |
| Mallow | | | | |
| Common (Cheeseweed) | <i>Malva neglecta</i> | | | |
| Little Mallow | <i>Malva parviflora</i> | | | |
| Horseweed/Marestail | <i>Conyza Canadensis</i> | | | |
| Morningglories | | | | |
| Entireleaf | <i>Ipomoea hederacea</i> var. <i>integriuscula</i> | | | |
| Ivyleaf | <i>Ipomoea hederacea</i> | | | |
| Red/Scarlet | <i>Ipomoea coccinea</i> | | | |
| Smallflower | <i>Jacquemontia tamnifolia</i> | | | |
| Tall | <i>Ipomoea purpurea</i> | | | |

continued

¹ *Chateau* WDG can be used on soils with greater than 10% organic matter; however, length of residual control may be shorter than on soils with lower organic matter content.
² A maximum *Chateau* WDG rate of 6 oz/A per application should be used on any soil that has a sand plus gravel content over 80% if trees are under 3 years of age.

Table 1. Weeds Controlled by Preemergence Application of *Chateau* WDG (continued)

| BROADLEAF WEED SPECIES | | | | |
|-------------------------------|--|------------------------|-----------------------------|---------------------------|
| COMMON NAME | SCIENTIFIC NAME | ORGANIC MATTER | SOIL TYPE | CHATEAU WDG RATE |
| Mustards | | Up to 10% ¹ | All Soil Types ² | 6 to 12 oz/A ² |
| Tumble | <i>Sisymbrium altissimum</i> | | | |
| Wild | <i>Brassica kaber</i> | | | |
| Nightshades | | | | |
| Black | <i>Solanum nigrum</i> | | | |
| Eastern Black | <i>Solanum ptycanthum</i> | | | |
| Hairy | <i>Solanum sarrachoides</i> | | | |
| Pigweeds | | | | |
| Palmer Amaranth | <i>Amaranthus palmeri</i> | | | |
| Redroot | <i>Amaranthus retroflexus</i> | | | |
| Smooth | <i>Amaranthus hybridus</i> | | | |
| Spiny Amaranth | <i>Amaranthus spinosus</i> | | | |
| Tumble | <i>Amaranthus albus</i> | | | |
| Prickly Sida (Teaweed) | <i>Sida spinosa</i> | | | |
| Puncturevine | <i>Tribulus terrestris</i> | | | |
| Purslane, Common | <i>Portulaca oleracea</i> | | | |
| Radish, Wild | <i>Raphanus raphanistrum</i> | | | |
| Ragweed, Common | <i>Ambrosia artemisiifolia</i> | | | |
| Redmaids | <i>Calandrinia ciliata</i> var. <i>menziessi</i> | | | |
| Redweed | <i>Melochia corchorifolia</i> | | | |
| Shepherd's-Purse | <i>Capsella bursa-pastoris</i> | | | |
| Sowthistle, Annual | <i>Sonchus oleraceus</i> | | | |
| Spotted Spurge | <i>Euphorbia maculata</i> | | | |
| Spurred Anoda | <i>Anoda cristata</i> | | | |
| Thistle, Russian | <i>Salsola iberica</i> | | | |
| Tropic Croton | <i>Croton glandulosus</i> | | | |
| Venice Mallow | <i>Hibiscus trionum</i> | | | |
| Waterhemp | | | | |
| Common | <i>Amaranthus rudis</i> | | | |
| Tall | <i>Amaranthus tuberculatus</i> | | | |
| Wild Poinsettia | <i>Euphorbia heterophylla</i> | | | |

continued

¹ *Chateau* WDG can be used on soils with greater than 10% organic matter; however, length of residual control may be shorter than on soils with lower organic matter content.

² A maximum *Chateau* WDG rate of 6 oz/A per application should be used on any soil that has a sand plus gravel content over 80% if trees are under 3 years of age.

Table 1. Weeds Controlled by Preemergence Application of *Chateau* WDG (continued)

| GRASS WEED SPECIES | | | | |
|---------------------------|--------------------------------|-----------------------|-----------------------------|---------------------------|
| COMMON NAME | SCIENTIFIC NAME | ORGANIC MATTER | SOIL TYPE | CHATEAU WDG RATE |
| Barnyardgrass | <i>Echinochloa crus-galli</i> | Up to 10% | All Soil Types ² | 6 to 12 oz/A ² |
| Bluegrass, Annual | <i>Poa annua</i> | | | |
| California Lovegrass | <i>Eragrostis diffusa</i> | | | |
| Crabgrass | | | | |
| Large | <i>Digitaria sanguinalis</i> | | | |
| Smooth | <i>Digitaria ischaemum</i> | | | |
| Foxtails | | | | |
| Bristly | <i>Setaria verticillata</i> | | | |
| Giant | <i>Setaria faberi</i> | | | |
| Green | <i>Setaria viridis</i> | | | |
| Yellow | <i>Setaria glauca</i> | | | |
| Goosegrass | <i>Eleusine indica</i> | | | |
| Guineagrass | <i>Panicum maximum</i> | | | |
| Johnsongrass, Seedling | <i>Sorghum halepense</i> | | | |
| Signalgrass, Broadleaf | <i>Brachiaria platyphylla</i> | | | |
| Panicum | | | | |
| Fall | <i>Panicum dichotomiflorum</i> | | | |
| Texas | <i>Panicum texaum</i> | | | |

¹ *Chateau* WDG can be used on soils with greater than 10% organic matter; however, length of residual control may be shorter than on soils with lower organic matter content.

² A maximum *Chateau* WDG rate of 6 oz/A per application should be used on any soil that has a sand plus gravel content over 80% if trees are under 3 years of age.

Table 2. Weeds Controlled by Postemergence Activity of *Chateau* WDG Tank Mixes

| BROADLEAF WEED SPECIES | | | |
|---------------------------------------|--|------------------------------------|-------------------------|
| COMMON NAME | SCIENTIFIC NAME | WEED HEIGHT/LENGTH (inches) | CHATEAU WDG RATE |
| Bindweed, Field ¹ | <i>Convolvulus arvensis</i> | 8 | 6 to 12 oz/A |
| Carpetweed | <i>Mollugo verticillata</i> | 4 | |
| Chickweeds | | | |
| Common | <i>Stellaria media</i> | 4 | |
| Mouseear | <i>Cerastium vulgatum</i> | 4 | |
| Cocklebur, Common | <i>Xanthium strumarium</i> | 4 | |
| Eveningprimrose, Cutleaf ² | <i>Oenothera laciniata</i> | 12 | |
| Filaree | | | |
| Broadleaf | <i>Erodium botrys</i> | 4 | |
| Redstem | <i>Erodium cicutarium</i> | 4 | |
| Florida Beggarweed | <i>Desmodium tortuosum</i> | 2 | |
| Hemp Sesbania | <i>Sesbania exaltata</i> | 8 | |
| Jimsonweed | <i>Datura stramonium</i> | 4 | |
| Lambsquarters, Common | <i>Chenopodium album</i> | 4 | |
| Morningglories | | | |
| Entireleaf | <i>Ipomoea hederacea</i> var. <i>integriuscula</i> | 4 | |
| Ivyleaf | <i>Ipomoea hederacea</i> | 4 | |
| Pitted | <i>Ipomoea lacunosa</i> | 6 | |
| Red/Scarlet | <i>Ipomoea coccinea</i> | 4 | |
| Tall | <i>Ipomoea purpurea</i> | 4 | |
| Pigweeds | | | |
| Palmer Amaranth | <i>Amaranthus palmeri</i> | 6 | |
| Redroot | <i>Amaranthus retroflexus</i> | 6 | |
| Smooth | <i>Amaranthus hybridus</i> | 6 | |
| Plantain, Broadleaf | <i>Plantago major</i> | 6 | |
| Prickly Sida (Teaweed) | <i>Sida spinosa</i> | 6 | |
| Purslanes | | | |
| Common | <i>Portulaca oleracea</i> | 4 | |
| Rock | <i>Calandrinia</i> spp. | 2 | |
| Ragweeds | | | |
| Common | <i>Ambrosia artemisiifolia</i> | 2 | |
| Giant | <i>Ambrosia trifida</i> | 4 | |
| Sicklepod | <i>Senna obtusifolia</i> | 4 | |

continued

¹ *Chateau* WDG will only provide control of the above ground portion of bindweed. Repeated applications will be needed to control regrowth.

² For acceptable control, cutleaf eveningprimrose should be 12 inches or less and in the rosette stage. Crop oil concentrate, at 1 pt/A, or non-ionic surfactant at 0.25% v/v, should be added to glyphosate tank mixes for cutleaf eveningprimrose control, including glyphosate formulations that contain a built-in adjuvant system.

Table 2. Weeds Controlled by Postemergence Activity of *Chateau* WDG Tank Mixes
(continued)

| BROADLEAF WEED SPECIES | | | |
|-------------------------------|---------------------------------|------------------------------------|-------------------------|
| COMMON NAME | SCIENTIFIC NAME | WEED HEIGHT/LENGTH (inches) | CHATEAU WDG RATE |
| Smartweeds | | | 6 to 12 oz/A |
| Ladysthumb | <i>Polygonum persicaria</i> | 4 | |
| Pale | <i>Polygonum lapathifolium</i> | 4 | |
| Pennsylvania | <i>Polygonum pennsylvanicum</i> | 4 | |
| Rice Flatsedge | <i>Cyperus iria</i> | 4 | |
| Spotted Spurge | <i>Euphorbia maculata</i> | 4 | |
| Velvetleaf | <i>Abutilon theophrasti</i> | 4 | |
| Venice Mallow | <i>Hibiscus trionum</i> | 4 | |
| Waterhemp | | | |
| Common | <i>Amaranthus rudis</i> | 2 | |
| Tall | <i>Amaranthus tuberculatus</i> | 2 | |
| Wild Mustard | <i>Brassica kaber</i> | 6 | |

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. PLEASE REFER TO CONTAINER LABEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS. FOLLOW ALL APPLICATION DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED LABEL.

PLEASE CONTACT VALENT U.S.A. CORPORATION AT 800-6-VALENT (682-5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE.

Copyright © 2008 by Valent U.S.A. Corporation

Chateau and *Products That Work, From People Who Care* are registered trademarks of Valent U.S.A. Corporation

Rely is a registered trademark of Bayer

Surflan is a registered trademark of United Phosphorus, Inc.

Manufactured for
Valent U.S.A. Corporation
P. O. Box 8025
Walnut Creek, CA 94596-8025

Made in U.S.A.

Form 2008-CHA-0029
(FPL - 20059639.00119.20080909.CHAWDG.FINAL)

Products That Work, From People Who Care® | www.valent.com | 800-6-VALENT (682-5368)