

# Supplemental Label



EPA Reg. No. 59639-99  
(For California Only)

## CHATEAU® HERBICIDE SW USE IN STRAWBERRY

**THIS SUPPLEMENTAL LABEL SURPERCEDES PREVIOUS LABELING FOR USE IN STRAWBERRY.**

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.**

### GENERAL RESTRICTIONS AND LIMITATIONS:

- Do not apply more than 3 oz of *Chateau* SW per acre per application.
- Do not apply more than 3 oz of *Chateau* SW per acre during a single growing season.
- *Chateau* SW, at 3 oz per acre, can be applied to the soil a minimum of 30 days prior to transplanting strawberries provided the strawberries will be transplanted through a plastic mulch.
- *Chateau* SW, at 3 oz per acre, can be applied in strawberry row middles with a shielded or hooded sprayer for the preemergence control of the weeds listed in Table 1, Broadleaf Weeds Controlled by Residual Activity of *Chateau* SW.

Application Method	Minimum Time From Application to Harvest (PHI)	Use Rate Per Acre Per Application (oz)	Use Rate Per Acre Per Year (oz)	Special Use Instructions
Pre-transplant	Not applicable	3	3	Apply a minimum of 30 days prior to transplanting and prior to plastic mulch being laid.  Apply as part of a tank mix to control emerged weeds.

continued

Application Method	Minimum Time From Application to Harvest (PHI)	Use Rate Per Acre Per Application (oz)	Use Rate Per Acre Per Year (oz)	Special Use Instructions
Hooded or shielded sprayer application to row middles	Do not apply after fruit set	3	3	<p><b>Apply only to row middles - do not apply over strawberries.</b></p> <p>Apply prior to weed emergence.</p> <p>Crop spotting may occur if an adjuvant is added.</p> <p><b>Application after fruit set may result in spotting of fruit and should be avoided.</b></p> <p>Do not allow spray drift to come in contact with fruit or foliage</p>

**Table 1. Broadleaf Weeds Controlled by Residual Activity of *Chateau* SW**

BROADLEAF WEED SPECIES					
SECTION A					
COMMON NAME	SCIENTIFIC NAME	ORGANIC MATTER	SOIL TYPE	CHATEAU SW RATE	
Carpetweed	<i>Mollugo verticillata</i>	Up to 5%	All Soil Types	2 oz/A	
Chickweeds					
Common	<i>Stellaria media</i>				
Mouseear	<i>Cerastium vulgatum</i>				
Dandelion	<i>Taraxacum officinale</i>				
Eclipta	<i>Eclipta prostrata</i>				
Eveningprimrose, Cutleaf	<i>Oenothera laciniata</i>				
Florida Pusley	<i>Richardia scabra</i>				
Henbit	<i>Lamium amplexicaule</i>				
Lambsquarters, Common	<i>Chenopodium album</i>				
Little Mallow	<i>Malva parviflora</i>				
Marestail/Horseweed	<i>Conyza canadensis</i>				
Nightshades					
Black	<i>Solanum nigrum</i>				
Eastern Black	<i>Solanum ptycanthum</i>				
Hairy	<i>Solanum sarrachoides</i>				
Pigweeds					
Redroot	<i>Amaranthus retroflexus</i>				
Smooth	<i>Amaranthus hybridus</i>				
Spiny Amaranth	<i>Amaranthus spinosus</i>				
Tumble	<i>Amaranthus albus</i>				
Prickly Sida (Teaweed)	<i>Sida spinosa</i>				
Puncturevine	<i>Tribulus terrestris</i>				
Purslane, Common	<i>Portulaca oleracea</i>				
Radish, Wild	<i>Raphanus raphanistrum</i>				
Redmaids	<i>Calandrinia ciliata var menziessii</i>				
Shepherd's-purse	<i>Capsella bursa-pastoris</i>				
Smallflower Morningglory	<i>Jacquemontia tamnifolia</i>				
Spotted Spurge	<i>Euphorbia maculata</i>				
Venice Mallow	<i>Hibiscus trionum</i>				

continued

**Table 1. Broadleaf Weeds Controlled by Residual Activity of *Chateau SW* (continued)**

<b>SECTION B</b>				
<b>All weeds listed in Section A plus:</b>				
<b>COMMON NAME</b>	<b>SCIENTIFIC NAME</b>	<b>ORGANIC MATTER</b>	<b>SOIL TYPE</b>	<b>CHATEAU SW RATE</b>
Coffee Senna	<i>Cassia occidentalis</i>	Up to 3%	All Soil Types	3 oz/A
Common Ragweed	<i>Ambrosia artemisiifolia</i>			
Florida Beggarweed	<i>Desmodium tortuosum</i>			
Golden Crownbeard	<i>Verbesina encelioides</i>			
Hairy Indigo	<i>Indigofera hirsuta</i>			
Hemp Sesbania	<i>Sesbania exaltata</i>	3 to 5%	Coarse and Medium Soils: (sandy loam, loamy sand, loamy, silt-loam, silt, sandy clay, sandy clay loam)	3 oz/A
Kochia	<i>Kochia scoparia</i>			
Jimsonweed	<i>Datura stramonium</i>			
Morningglories <sup>1</sup>				
Entireleaf	<i>Ipomoea hederacea var integruscula</i>			
Ivyleaf	<i>Ipomoea hederacea</i>			
Red/Scarlet	<i>Ipomoea coccinea</i>			
Tall	<i>Ipomoea purpurea</i>			
Mustard, Wild	<i>Brassica kaber</i>			
Palmer Amaranth	<i>Amaranthus palmeri</i>			
Spurred Anoda	<i>Anoda cristata</i>		Fine Soils: (silty clay, silty clay loam, clay, clay loam)	3 oz/A
Tropic Croton	<i>Croton glandulosus</i>			
Waterhemp				
Common	<i>Amaranthus rudis</i>			
Tall	<i>Amaranthus tuberculatus</i>			
Wild Poinsettia	<i>Euphorbia heterophylla</i>			

<sup>1</sup>Morningglory species are not adequately controlled on fine soils or soils with greater than 3% organic matter.

**THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. PLEASE REFER TO CONTAINER LABEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS. FOLLOW ALL APPLICATION DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL. PLEASE CONTACT VALENT U.S.A. CORPORATION AT 800-6-VALENT (682-5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE.**

Copyright © 2008 by Valent U.S.A. Corporation

*Chateau* and *Products That work, From People Who Care* are registered trademarks of Valent U.S.A. Corporation

Manufactured for  
Valent U.S.A. Corporation  
P.O. Box 8025  
Walnut Creek, CA 94596-8025  
www.valent.com

Made in U.S.A.

Form 2008-CHSW-0034-CA

*Products That Work, From People Who Care*® | [www.valent.com](http://www.valent.com) | 800-6-VALENT (682-5368)