

Supplemental Label



PRESIDIO[®] FUNGICIDE

EPA Reg. No. 59639-140

PRESIDIO[®] FUNGICIDE FOR USE ON BRASSICA (HEAD AND STEM), BULB VEGETABLES AND ROOT VEGETABLES (EXCEPT CARROT, POTATO AND SUGAR BEET)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

ROTATIONAL RESTRICTIONS

The following rotational crops may be planted after applying *Presidio* Fungicide at the labeled rate. Planting earlier than the recommended rotational interval is not allowed.

CROPS	ROTATIONAL INTERVAL
Brassica (Head and Stem) Bulb Vegetables Cucurbit Vegetables Fruiting Vegetables Grapes Leafy Vegetables (Except Brassica) Root Vegetables, (Except Carrot, Potato and Sugar Beet)	Immediately
All Other Crops	18 months

SPECIFIC REQUIREMENTS – All Food Crops

When to Apply: Begin application when crop and/or environmental conditions favor disease development. Applications must be made on a 7 to 14 day interval, depending on disease pressure. Under severe disease pressure use the higher labeled rate and shortest interval. Apply as a foliar spray in sufficient water to obtain thorough coverage. Chemigation is not recommended when the primary disease is downy mildew or other foliar diseases.

continued

SPECIFIC REQUIREMENTS – All Food Crops (continued)

Resistance Management: *Presidio* Fungicide must be used as part of an integrated pest management (IPM) program. Apply as a foliar spray in a tank mix with another labeled fungicide product with a different mode of action on the target pathogen in sufficient water to obtain thorough coverage.

Restrictions and Limitations

- The maximum *Presidio* Fungicide application rate is 4 fl oz per acre per application and 12 fl oz per acre per season.
- Apply no more than 2 sequential applications of *Presidio* Fungicide before alternating with an effective fungicide from a different resistance management group.
- Do not make more than 4 applications of *Presidio* Fungicide per acre per season.

Crops	Diseases	Application Rates		Minimum Time from Last Application to Harvest (PHI)	Specific Use Instructions
		fl oz/A	GPA		
Brassica (Head and Stem): Broccoli Brussels Sprouts Cabbage Cauliflower Cavalo Broccolo Chinese Broccoli Chinese (napa) Cabbage Chinese Mustard Cabbage Kohlrabi	Downy Mildew (<i>Peronospora parasitica</i>) Phytophthora (<i>Phytophthora megasperma</i>)	3 to 4 (0.09 to 0.125 lb ai/A)	Ground: 20 to 50 Aerial Minimum: 5	2 days	A tank mix including a labeled rate of another labeled fungicide product with a different mode of action on the target pathogen must be used with <i>Presidio</i> Fungicide for resistance management. Recommended spray interval: 10 days

continued

continued

Crops	Diseases	Application Rates		Minimum Time from Last Application to Harvest (PHI)	Specific Use Instructions
		fl oz/A	GPA		
Bulb Vegetables including: Beltsville Bunching Onion Chive (fresh leaves) Chinese Chive (fresh leaves) Chinese Onion Bulb Daylily Bulb Elegans Hosta Fritillaria (bulb and leaves) Garlic Bulb Great Headed Garlic Bulb Green Onion Kurrant Lady's Leek Leek Lily Bulb Macrostem Onion Onion (bulb and fresh) Pearl Onion Potato Onion Bulb Shallot (bulb and fresh leaves) Serpent Garlic Bulb Tree Onion Tops Wild Leek	Downy Mildew (<i>Peronospora destructor</i>)	3 to 4 (0.09 to 0.125 lb ai/A)	Ground: 20 to 50 Aerial Minimum: 5	2 days	A tank mix including a labeled rate of another labeled fungicide product with a different mode of action on the target pathogen must be used with <i>Presidio</i> Fungicide for resistance management. Recommended spray interval: 10 days

continued

continued

Crops	Diseases	Application Rates		Minimum Time from Last Application to Harvest (PHI)	Specific Use Instructions
		fl oz/A	GPA		
Root Vegetables (Except Carrot, Potato and Sugar Beet): Arracacha Arrowroot Black Salsify Cassava (bitter and sweet) Celeriac Chayote (root) Chicory Chinese Artichoke Chufa Dasheen (taro) Edible Burdock Edible Canna Garden Beet Ginger Ginseng Jerusalem Artichoke Leren Horseradish Parsnip Radish, Oriental Radish Rutabaga Salsify Skirret Spanish Salsify Tanier True Yam Turmeric Turnip Turnip-rooted Chervil Turnip-rooted Parsley Yam Bean	Root Dieback (Forking and Stubbing) (<i>Pythium ultimum</i> and <i>Pythium irregulare</i>) Cavity Spot (<i>Pythium violae</i>)	3 to 4 (0.09 to 0.125 lb ai/A)	Ground: 20 to 50 Aerial Minimum: 5	7 days	A tank mix including a labeled rate of another labeled fungicide product with a different mode of action on the target pathogen must be used with <i>Presidio</i> Fungicide for resistance management. Recommended spray interval: 10 days

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. PLEASE REFER TO CONTAINER LABEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS. FOLLOW ALL APPLICATION DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL.

PLEASE CONTACT VALENT U.S.A. CORPORATION AT 800-6-VALENT (682-5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE.

Copyright © 2010 by Valent U.S.A. Corporation

Presidio and *Products That Work, From People Who Care* are registered trademarks of Valent U.S.A. Corporation

Manufactured for
Valent U.S.A. Corporation
P.O. Box 8025
Walnut Creek, CA 94596-8025

Form 2008-PRD-0011, R1

Products That Work, From People Who Care® | www.valent.com | 800-6-VALENT (682-5368)