

Supplemental Label



**RESTRICTED USE PESTICIDE
DUE TO TOXICITY TO FISH AND AQUATIC ORGANISMS.**

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.



EPA Reg. No. 59639-35

**DANITOL® 2.4 EC SPRAY FOR ADDITIONAL CROP USES ON BERRIES, OLIVE, POME FRUIT,
STONE FRUIT, TREE NUTS AND TROPICAL AND SUB-TROPICAL FRUIT**
**DANITOL 2.4 EC SPRAY FOR USE ON ADDITIONAL PESTS IN
FRUITING VEGETABLES, GRAPE AND PEA**
**DANITOL 2.4 EC SPRAY TANK MIX WITH SELECT MAX® HERBICIDE
WITH INSIDE TECHNOLOGY™ FOR USE ON COTTON**

**THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION.
READ THE LABEL AFFIXED TO THE CONTAINER FOR DANITOL 2.4 EC SPRAY BEFORE APPLYING.
USE OF DANITOL 2.4 EC SPRAY ACCORDING TO THIS LABELING IS SUBJECT TO THE USE
PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR
DANITOL 2.4 EC SPRAY**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

ADDITIONAL CROP USES

BERRIES

CROPS	PESTS	PRODUCT RATES FL OZ/A	SPECIAL INSTRUCTIONS
Bushberries: Blueberry Highbush Lowbush Rabbiteye Elderberry Gooseberry Huckleberry Juneberry	Blueberry Maggot Cherry Fruitworm Cranberry Fruitworm Japanese Beetle Obliquebanded Leafroller Plum Curculio	10-2/3 to 16 (0.2 to 0.3 lb ai/A)	Apply with air or ground equipment in adequate water for uniform coverage (3 to 10 gals/A by air or a minimum of 20 gals/A by ground). Make applications beginning when first pest activity is noticed and repeat as needed but not more often than every 14 days.

continued

BERRIES (continued)

CROPS	PESTS	PRODUCT RATES FL OZ/A	SPECIAL INSTRUCTIONS
Lingonberry Salal Caneberries: Blackberry (including Bingleberry Boysenberry Dewberry Lowberry Marionberry Olallieberry Youngberry) Loganberry Raspberry, Black and Red	European Red Mite Two-spotted Spider Mite	16 (0.3 lb ai/A)	Resistance Management: Do not make more than two (2) <i>Danitol</i> 2.4 EC Spray applications per season as part of a Resistance Management Program. Use non-pyrethroid products at other timings to control pests.
	Blueberry Maggot Cherry Fruitworm Cranberry Fruitworm Japanese Beetle Obliquebanded Leafroller Plum Curculio	10-2/3 to 16 (0.2 to 0.3 lb ai/A)	
	European Red Mite Two-spotted Spider Mite	16 (0.3 lb ai/A)	
<ul style="list-style-type: none"> Do not apply within 3 days of harvest. Do not exceed 2 pts (32 fl oz, 0.6 lb ai) total application of <i>Danitol</i> 2.4 EC Spray per acre per season. 			
Currant	Currant Borer	10-2/3 to 16 (0.2 to 0.3 lb ai/A)	Apply as a ground application in a minimum of 50 gals of water per acre. Make applications beginning when first pest activity is noticed and repeat as needed but not more often than every 7 days. Resistance Management: Do not make more than three (3) <i>Danitol</i> 2.4 EC Spray applications per season as part of a Resistance Management Program. Use non-pyrethroid products at other timings to control pests.
<ul style="list-style-type: none"> Do not apply within 21 days of harvest. Do not exceed 2-2/3 pts (42-2/3 fl oz, 0.8 lb ai) total application of <i>Danitol</i> 2.4 EC Spray per acre per season. 			

OLIVE

PEST	PRODUCT RATE FL OZ/A	SPECIAL INSTRUCTIONS
Olive Fruit Fly	10-2/3 to 16 (0.2 to 0.3 lb ai/A)	<p>Apply as a ground application in a minimum of 100 gallons of water per acre.</p> <p>Make applications beginning when first pest activity is noticed and repeat as needed but not more often than every 14 days.</p> <p>Under severe pest pressure use the higher recommended rate.</p>
<ul style="list-style-type: none"> Do not apply within seven (7) days of harvest. Do not exceed 2-2/3 pts (42-2/3 fl oz, 0.8 lb ai) total application of <i>Danitol</i> 2.4 EC Spray per acre per season. 		

POME FRUIT: Apple, Crabapple, Loquat, Mayhaw, Quince

PESTS	PRODUCT RATES FL OZ/A	SPECIAL INSTRUCTIONS
Potato Leafhopper Rosy Apple Aphid Spotted Tentiform Leafminer Tarnished Plant Bug White Apple Leafhopper	10-2/3 to 21-1/3 (0.2 to 0.4 lb ai/A)	<p>Begin applications at delayed dormant through first cover as common to the production area and the target pest.</p> <p>Apply by ground with airblast equipment as a full coverage spray and repeat as needed to maintain control but not more often than every 10 days.</p>
Apple Maggot Codling Moth European Apple Sawfly European Red Mite Green Fruitworm Green Stinkbug Japanese Beetle Lesser Appleworm Lygus Bugs Obliquebanded Leafroller Oriental Fruitmoth Pandemis Leafroller Periodical Cicada Plum Curculio Redbanded Leafroller Spirea Aphid Southern Green Stinkbugs Tufted Apple Budmoth Two-spotted Spider Mite Variegated Leafroller	16 to 21-1/3 (0.3 to 0.4lb ai/A)	<p>Resistance Management: It is generally recommended that no more than 2 <i>Danitol</i> 2.4 EC Spray applications be made per season as part of a Resistance Management Program. Use non-pyrethroid products at other timings to control pests.</p> <p>Aphid Control: Timing of applications is critical in achieving control. Use prebloom spray timings for Rosy Apple Aphid and post bloom spray timings for Spirea Aphid as recommended by State or County Extension Service.</p> <p>Codling Moth: Make first application at Biofix + 250 degree days and repeat as needed to maintain control but not more often than every 10 days.</p> <p>Make pyrethroid insecticide applications according to local IPM programs and guidelines.</p> <p>Overwintering Pandemis and Obliquebanded leafrollers: Apply at pink stage of tree growth.</p>
<ul style="list-style-type: none"> Do not apply within 14 days of harvest. Do not exceed 2-2/3 pts (42-2/3 fl oz, 0.8 lb ai) total application of <i>Danitol</i> 2.4 EC Spray per acre per season Do not apply as a ULV spray. Do not feed or allow livestock to graze on cover crops from treated orchards. 		

STONE FRUIT: such as Apricot, Cherry, Nectarine, Peach, Plum, Prune

PESTS	PRODUCT RATE FL OZ/A	SPECIAL INSTRUCTIONS
European Red Mite Green Fruitworm Green Stinkbug Navel Orangeworm Obliquebanded Leafroller Orange Tortrix Oriental Fruitmoth Peach Twig Borer Redbanded Leafroller Southern Green Stinkbug Two-spotted Spider Mite Variegated Leafroller	10-2/3 to 21-1/3 (0.2 to 0.4 lb ai/A)	Apply by ground in 100 to 400 gals/A with airblast equipment as a full coverage spray and repeat as needed to maintain control but not more often than every ten (10) days. Coverage is essential for good control. Apply in sufficient water volume for tree size. Use of higher water volume will assure better coverage. Begin applications as common to the production area and the target pest. Resistance Management: It is generally recommended that no more than two (2) <i>Danitol 2.4 EC Spray</i> applications be made per season as part of a Resistance Management program. Use non-pyrethroid products at other timings to control pests. Make pyrethroid insecticide applications according to local Integrated Pest Management (IPM) programs and guidelines.
<ul style="list-style-type: none"> Do not apply <i>Danitol 2.4 EC Spray</i> within 3 days of harvest. Do not exceed 2-2/3 pts (42-2/3 fl oz, 0.8 lb ai) total application of <i>Danitol 2.4 EC Spray</i> per acre per season. Do not apply as an Ultra Low Volume (ULV) spray. Do not feed or allow livestock to graze on cover crops from treated orchards. 		

TREE NUTS: such as Almond, Beechnut, Brazil, Butternut, Cashew, Chestnut, Chinquapin, Filbert (Hazelnut), Hickory, Macadamia, Pecan, Pistachio, Walnut

PESTS	PRODUCT RATE FL OZ/A	SPECIAL INSTRUCTIONS
Codling Moth European Red Mite Green Fruitworm Green Stinkbug Leafroller Navel Orangeworm Obliquebanded Leafroller Twig Borer Pecan Nut Casebearer Plant Bug Redbanded Leafroller Shuckworm Southern Green Stinkbug Two-spotted Spider Mite Variegated Leafroller	10-2/3 to 21-1/3 (0.2 to 0.4 lb ai/A)	Apply by ground in 50 to 400 gals/A with airblast equipment as a full coverage spray and repeat as needed to maintain control but not more often than every 10 days. Coverage is essential for good control. Apply in sufficient water volume for tree size. Use of higher water volume will assure better coverage. Begin applications as common to the production area and the target pest. Codling Moth: Make first application at Biofix + 250 degree days and repeat as needed to maintain control but not more often than every 10 days. Resistance Management: It is generally recommended that no more than two (2) <i>Danitol 2.4 EC Spray</i> applications be made per season as part of a Resistance Management program. Use non-pyrethroid products at other timings to control pests. Make pyrethroid insecticide applications according to local Integrated Pest Management (IPM) programs and guidelines.

continued

TREE NUTS: such as Almond, Beechnut, Brazil, Butternut, Cashew, Chestnut, Chinquapin, Filbert (Hazelnut), Hickory, Macadamia, Pecan, Pistachio, Walnut – continued

- Do not apply *Danitol* 2.4 EC Spray within 3 days of harvest.
- Do not exceed 2-2/3 pts (42-2/3 fl oz, 0.8 lb ai) total application of *Danitol* 2.4 EC Spray per acre per season.
- Do not apply as an Ultra Low Volume (ULV) spray.
- Do not feed or allow livestock to graze on cover crops from treated orchards.

TROPICAL AND SUB TROPICAL FRUIT (Inedible Peel): Avocado, Canistel, Mango, Papaya, Sapodilla, Black Sapote, Mamey Sapote, Star Apple

PESTS	PRODUCT RATES FL OZ/A	SPECIAL INSTRUCTIONS
Avocado Looper Avocado Thrips Brown Mite Mirads Persea Mite	16 to 21-1/3 (0.3 to 0.4 lb ai/A)	Apply as a ground application in a minimum of 100 gals of water per acre. Application to avocado may be with aircraft in a minimum of 50 gallons of water per acre. Make applications beginning when first pest activity is noticed and repeat as needed but not more often than every 14 days. Resistance Management: Avocado - Do not make more than one pyrethroid application per season as part of a Resistance Management Program. Other crops - Do not make more than two (2) <i>Danitol</i> 2.4 EC Spray applications per season as part of a Resistance Management Program. Use non-pyrethroid products at other timings to control pests.
<ul style="list-style-type: none"> • Do not exceed 2-2/3 pts (42-2/3 fl oz, 0.8 lb ai) of <i>Danitol</i> 2.4 EC Spray per acre per season. • Do not apply within 1 day of harvest. 		

DANITOL 2.4 EC SPRAY FOR USE ON ADDITIONAL PESTS

FRUITING VEGETABLES: Bell Pepper, Chili Pepper, Cooking Pepper, Eggplant, Ground Cherry, (*Physalis* spp.), Pepino, Pimento, Sweet Pepper, Tomatillo, Tomato

PESTS	PRODUCT RATE FL OZ/A	SPECIAL INSTRUCTIONS
Green Stinkbug Southern Green Stinkbug See container label for more pests.	10-2/3 (0.2 lb ai/A)	See container label for special instructions and precautions.

GRAPE

PESTS	PRODUCT RATE FL OZ/A	SPECIAL INSTRUCTIONS
Plum Curculio See container label for more pests.	16 to 21-1/3 (0.3 to 0.4lb ai/A)	See container label for special instructions and precautions.

PEA (Succulent) including: English Pea, Garden Pea, Green Pea, Pigeon Pea

PESTS	PRODUCT RATE FL OZ/A	SPECIAL INSTRUCTIONS
Green Stinkbug Southern Green Stinkbug See container label for more pests.	10-2/3 (0.2 lb ai/A)	See container label for special instructions and precautions.

**TANK MIX APPLICATION OF *DANITOL* 2.4 EC SPRAY INSECTICIDE/MITICIDE AND *SELECT MAX*
HERBICIDE WITH *INSIDE TECHNOLOGY* FOR CONTROL OF INSECTS/MITES AND GRASS
FOR USE ON COTTON**

APPLICATION RATES/ACRE ¹			
PRODUCT	ANNUAL GRASSES	PERENNIAL GRASSES	ADJUVANT RATE (V/V) ²
<i>Danitol</i> 2.4 EC Spray + <i>Select Max</i> Herbicide with <i>Inside Technology</i>	10-2/3 to 16 fl oz + 9 to 16 fl oz	10-2/3 to 16 fl oz + 12 to 32 fl oz	See tank mix label for specific adjuvant recommendations

¹ If grass regrowth occurs, or an additional flush of new grass emerges, make a second application of *Select Max* Herbicide with *Inside Technology* alone (without *Danitol* 2.4 EC Spray) according to the appropriate size and rate recommendations.

² Always use non-ionic surfactant at 0.25% or COC/MSO at 1 qt/A or 1% v/v in the finished spray solution plus ammonium sulfate at 2.5 to 4.0 lbs/A.

PLEASE CONTACT VALENT U.S.A. CORPORATION AT 800-6-VALENT (682-5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE.

Copyright© 2009 by Valent U.S.A. Corporation

with Inside Technology is a trademark and *Select Max* and Products That Work, From People Who Care are registered trademarks of Valent U.S.A. Corporation. *Danitol* is a registered trademark of Sumitomo Chemical Company, Ltd.

Manufactured for
Valent U.S.A. Corporation
P. O. Box 8025
Walnut Creek, CA 94596-8025
Made in U.S.A.

Form 2009-DAN-0011R1

Products That Work, From People Who Care® | www.valent.com | 800-6-VALENT (682-5368)