



**Active Ingredient<sup>1</sup>:**  
*Myrothecium verrucaria* strain AARC-0255  
 fermentation solids and solubles ..... 90% w/w  
**Other Ingredients** ..... 10% w/w  
**Total** ..... 100% w/w  
<sup>1</sup>“Non-viable”/“killed” microbial composition

Potency: 91,600 RKU (Root-Knot Units) per gram of product.  
**Potency units should not be used to adjust use rates.**

EPA Reg. No. 73049-67  
 EPA Est. No. 33762-IA-001  
 List No. 60278

**INDEX:**

- 1.0 First Aid
- 2.0 Precautionary Statements
  - 2.1 Hazards to Humans and Domestic Animals
  - 2.2 Personal Protective Equipment (PPE)
  - 2.3 User Safety Recommendations
  - 2.4 Environmental Hazards
- 3.0 Directions for Use
- 4.0 Agricultural Use Requirements
- 5.0 Non-Agricultural Use Requirements
- 6.0 Storage and Disposal
- 7.0 General Information
- 8.0 General Instructions
- 9.0 Compatibility
- 10.0 Chemigation
  - 10.1 Mixing and Application Instructions
  - 10.2 Chemigation Use Directions
- 11.0 Table 1
- 12.0 Table 2. Conversion Table for Band Applications
- 13.0 Notice to User

**KEEP OUT OF REACH OF CHILDREN  
 CAUTION**

1.0 FIRST AID	
<b>If in Eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on Skin or Clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergencies, you may also call toll-free 1-800-892-0099 for treatment information.	

**2.0 PRECAUTIONARY STATEMENTS**

**2.1 HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
 CAUTION**

Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

**2.2 Personal Protective Equipment (PPE)**

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow manufacturer’s instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**2.3 User Safety Recommendations**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

**2.4 Environmental Hazards**

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

**3.0 DIRECTIONS FOR USE**

It is a violation to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system unless labeling for chemigation is followed.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the Agency responsible for pesticide regulations.

#### 4.0 AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

Exception: If the product is soil injected or soil incorporated, the Worker Protection Standard under certain circumstances allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

#### 5.0 NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries.

Keep unprotected persons out of treated areas until sprays have dried.

#### 6.0 STORAGE AND DISPOSAL

Do not contaminate water, food or feed by improper storage or disposal.

**Pesticide Storage:** Keep containers tightly closed when not in use. Store in a cool, dry place. Avoid extreme temperatures.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Do not contaminate water when disposing of equipment washwaters.

**Container Disposal:** Non refillable container. Do not reuse or refill this container. Offer for recycling, if available. Completely empty container into applicable equipment. If recycling is not available, dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

#### 7.0 GENERAL INFORMATION

*DiTera* DF controls the following plant nematodes parasitizing food, fiber and ornamental crops: *Meloidogyne* spp. (root-knot nematodes), *Heterodera* and *Globodera* spp. (cyst nematodes), *Pratylenchus* spp. (lesion nematodes), *Tylenchulus semipenetrans* (citrus nematode), *Trichodorus* spp. (stubby-root nematodes), *Longidorus* spp. (needle nematodes), *Paratylenchus* spp. (pin nematodes), *Rotylenchulus* spp. (reniform nematodes), *Xiphinema* spp. (dagger nematodes), *Belonolaimus* spp. (sting nematodes), *Criconeoides* spp., *Criconebella* spp. and related genera (ring nematodes), *Tylenchorhynchus* spp. (stunt nematodes), *Hoplolaimus* spp. (lance nematodes), *Rotylenchus* spp., *Helicotylenchus* spp. (spiral nematodes), *Radopholus* spp. (burrowing nematodes) and other plant parasitic nematodes.

#### 8.0 GENERAL INSTRUCTIONS

*DiTera* DF may be applied to the soil as a pre-plant, at planting or post-plant treatment on annual and perennial crops using ground equipment (refer to commodities listed in Table 1). Pre-plant applications should preferably be close to the actual planting times. The optimal application time should be determined based on the cultural practices and the nematode population dynamics. For perennial crops, apply *DiTera* DF just prior to a root flush to protect young roots. Multiple applications may be required for crops with multiple root flushes. *DiTera* DF must be incorporated into the soil for optimum results. Incorporation may be accomplished by mechanical equipment, irrigation or rainfall. For applications made at planting, the action of some planters may provide sufficient incorporation. When using planters which do not provide adequate incorporation of *DiTera* DF into the soil, equipment designed for incorporation may be used behind the planter.

The rates provided are for broadcast applications. Due to the nature of the active ingredient and the distribution of nematodes in agricultural soils, in-row or band application may be suitable for adequate nematode suppression. If banding or side-dressing, the corresponding rates need to be calculated based on the actual surface area of soil to be treated (Refer to conversion table for band applications). Use of the higher rates and/or multiple applications may be required in coarse (light) soils with less than 1% organic matter. Maximum benefits may not be realized in agricultural fields containing non-decomposed plant materials, including infected roots from a previous crop, or in fields with very high nematode infestations.

*DiTera* DF may be spread as a dry granule, applied using ground sprays, or using approved irrigation systems (refer to Chemigation Use Directions), with quantities of water sufficient to provide coverage of the root area of the plants. The amount of water needed per acre will depend on the plant species, biology of the nematode species to be controlled, stage of crop, weather, soil moisture conditions, level of nematode infestation, etc. Do not apply by mist sprayer or aerial spray equipment.

**9.0 COMPATIBILITY**

The *DiTera* DF application directions refer to the use of the product alone. Data concerning the compatibility of *DiTera* DF with other agricultural products are not available. Valent does not assume responsibility for unexpected, adverse results due to the tank mixing or simultaneous applications of *DiTera* DF with other agricultural products including fertilizers.

**10.0 CHEMIGATION**

**10.1 MIXING AND APPLICATION INSTRUCTIONS**

To ensure complete mixing of *DiTera* DF, spray equipment having an agitation system is recommended.

A thoroughly mixed *DiTera* DF suspension can be applied through irrigation systems described under Chemigation Use Directions, so as to treat the soil around the crop root zone. It is recommended that the material be used immediately after mixing in order to avoid settling. *DiTera* DF must be applied as a dedicated irrigation between normal irrigations or at the end of a scheduled irrigation. To minimize excessive dilution or leaching of product from root zone, *DiTera* DF must be applied in the highest concentration over the shortest time span possible.

Pre-mix product in a mix tank having both mechanical and by-pass agitation. Fill tank with 1/2 to 3/4 of the desired amount of water. Start mechanical and hydraulic systems to provide maximum agitation. Add an agricultural wetting agent prior to adding *DiTera* DF. Add 1/3 to 1/2 of *DiTera* DF and allow to mix until thoroughly suspended. Add the remaining material slowly while bringing water to the desired volume. Continue agitation. If a concentration greater than 1 lb. *DiTera* DF per 2 gallons of water is desired, the material should be added in smaller increments with sufficient agitation following each addition. To avoid excessive leaching of product, flush irrigation lines for the minimum time period at the end of the application. Do not mix more *DiTera* DF than can be used in a 24-hour period. Rinse and flush spray equipment thoroughly following each use. Use a strainer no finer than 50 mesh in conventional spray systems.

**10.2 CHEMIGATION USE DIRECTIONS**

Apply this product only through the following types of irrigation systems: sprinkler, including center pivot, lateral move, end tow, side (wheel) roll, traveler, solid set, or hand move; flood (basin); furrow; border; or low pressure (including drip/trickle, minisprinklers, drip tape, strip tubing and jets).

Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact local state extension personnel, equipment manufacturers or other experts.

Do not connect irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as a drop structure or weir box to decrease potential for water source contamination from backflow if water flow stops.

Systems utilizing sprinkler chemigation, a pressurized water and pesticide injection system, or drip/trickle chemigation, must meet the following requirements.

- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- For sprinkler chemigation, do not apply when wind speed favors drift beyond the area intended for treatment.

**11.0 TABLE 1**

Crop Groups/ Representative Commodities	Nematode	Broadcast* Rate (pounds/ acre/appl)
<b>CITRUS FRUITS</b> such as sweet orange, lemon, grapefruit	Burrowing Citrus Cyst	13-100
<b>FLOWERING, BEDDING PLANTS, ORNAMENTALS,</b> such as fern and hosta	Dagger Lance Lesion	
<b>LEAFY VEGETABLES AND COLE CROPS</b> such as celery, head & leaf lettuce, spinach, broccoli, cabbage	Needle Pin Reniform	
<b>POME FRUITS</b> such as apple, pear	Ring Spiral Root-knot	
<b>STONE FRUITS</b> such as peach, plum, prune	Sting Stubby-root	
<b>TREE NUTS</b> such as almond, pecan, walnut	Stunt	
<b>GRAPE</b>		
<b>PINEAPPLE</b>		

\*Refer to accompanying table for band application rates.

CONTINUED

**12.0 TABLE 2 – CONVERSION TABLE  
FOR BAND APPLICATIONS**

Band Width Inches	Pounds of <i>DiTera</i> DF per 1,000 Feet of Row at Equivalent Broadcast Rate of Indicated Pounds per Acre			
	Equivalent Broadcast Rate Per Acre			
	13	25	50	100
12	0.31	0.60	1.2	2.4
18	0.47	0.91	1.8	3.6
24	0.63	1.20	2.4	4.8
30	0.79	1.51	3.0	6.0
36	0.94	1.82	3.6	7.2
48	1.26	2.40	4.8	9.6
60	1.58	3.00	6.0	12.0

**13.0 NOTICE TO USER**

Seller makes no warranty, express or implied, of merchantability, fitness or otherwise concerning the use of this product other than as indicated on the label. User assumes all risks of use, storage or handling not in strict accordance with accompanying directions.

*DiTera* is a registered trademark of Valent BioSciences Corporation.

*Products That Work, From People Who Care* is a registered trademark of Valent U.S.A. Corporation.

© 2010 Valent BioSciences Corporation

Registered and Manufactured by:  
Valent BioSciences Corporation  
870 Technology Way  
Libertyville, IL 60048



04-6542/R4 (Pkg. 04-6360/R4)

Distributed by Valent U.S. A. Corporation.  
P.O. Box 8025  
Walnut Creek, CA 94596-8025



For state registration and/or supplemental labels, please call or visit us online.

*Products That Work, From People Who Care*® | [www.valent.com](http://www.valent.com) | 800-6-VALENT (682-5368)

**Read and follow the label instructions before using.**

© 2010 Valent BioSciences Corporation. All rights reserved. Printed in the U.S.A. 2010-DITDF-0001 7/10 AV