

Supplemental Label



EPA Reg. No. 59639-99
(For Use in Colorado, Oregon and Washington Only)

VALOR[®] SX HERBICIDE USE ON DRY BEAN FOR WEED SUPPRESSION

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION.

READ THE LABEL AFFIXED TO THE CONTAINER FOR VALOR BEFORE APPLYING. USE OF VALOR ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR VALOR.

RESTRICTIONS AND LIMITATIONS

- Do not apply more than 1.5 oz of *Valor* per acre during a single application.
- Do not apply more than 1.5 oz of *Valor* per acre during a single growing season.

Many weather related factors, including high wind, splashing or heavy rains or cool conditions at or near crop emergence, may result in dry bean injury in fields treated with *Valor*. On occasion this has resulted in a delay in maturity. User should assume these risks before using *Valor*.

TIMING TO DRY BEAN

Valor may be applied to dry beans within 2 days after planting for the preemergence suppression of the weeds listed in Table 1, Weeds Suppressed by Residual Activity of *Valor*. *Valor* should be tank mixed with other labeled herbicides for broad spectrum weed control.

TIMING TO WEEDS

Valor may be applied to dry beans prior to planting or preemergence (after planting). Preemergence application of *Valor* must be made within 2 days after planting and prior to dry bean emergence. Application after the dry beans have begun to crack, or are emerged, will result in severe crop injury. Application should not be made when dry beans have begun to crack.

Preplant incorporation (PPI) applications may result in reduced weed control.

ADDITIONAL RESIDUAL GRASS CONTROL

Valor can be tank mixed with pendimethalin for additional grass control.

Read tank mix product label for rates and weeds controlled. Always read and follow label directions for all tank mix products before using. The most restrictive labeling of any tank mix product must be followed. *Valor*, when applied according to label use directions, will suppress the weeds listed in Table 1, Weeds Suppressed by Residual Activity of *Valor* at 1.5 oz/A. This label makes no claims concerning other weed species.

Table 1. Weeds Suppressed by Residual Activity of *Valor* at 1.5 oz/A

COMMON NAME	SCIENTIFIC NAME	ORGANIC MATTER	VALOR RATE
Lambsquarters, Common	<i>Chenopodium album</i>	Up to 5%	1.5 oz/A
Mustard, Wild	<i>Brassica kaber</i>		
Nightshades			
Black	<i>Solanum nigrum</i>		
Eastern Black	<i>Solanum ptycanthum</i>		
Hairy	<i>Solanum sarrachoides</i>		
Palmer Amaranth	<i>Amaranthus palmeri</i>		
Pigweeds			
Redroot	<i>Amaranthus retroflexus</i>		
Smooth	<i>Amaranthus hybridus</i>		
Spiny Amaranth	<i>Amaranthus spinosus</i>		
Tumble	<i>Amaranthus albus</i>		
Prickly Lettuce (China Lettuce)	<i>Lactuca serriola</i>		
Radish, Wild	<i>Raphanus raphanistrum</i>		

PLEASE CONTACT VALENT U.S.A. CORPORATION AT 800-6-VALENT (682-5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE.

Copyright © 2010 by Valent U.S.A. Corporation

Valor and *Products That Work, From People Who Care* are registered trademarks of Valent U.S.A. Corporation

Manufactured for
Valent U.S.A. Corporation
P. O. Box 8025
Walnut Creek, CA 94596-8025

Made in U.S.A.

Form 2010-VLR-0010

Products That Work, From People Who Care® | www.valent.com | 800-6-VALENT (682-5368)