

Supplemental Label



BELAY[®] **INSECTICIDE**

EPA Reg. No. 59639-150
(Except California, Florida and New York)

BELAY[®] INSECTICIDE FOR CRANBERRY, POTATO AND SWEET POTATO

This supplemental label expires on October 2, 2012 and must not be used or distributed after this date.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. READ THE LABEL AFFIXED TO THE CONTAINER FOR *BELAY* INSECTICIDE BEFORE APPLYING. USE OF *BELAY* INSECTICIDE ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR *BELAY* INSECTICIDE.

**CRANBERRY
(LOW-GROWING BERRY EXCEPT STRAWBERRY)**

Bearberry, Bilberry, Cloudberry, Cranberry, Lingonberry, Lowbush Blueberry, Muntries, Partridgeberry and varieties, and/or cultivars of these

FOLIAR POST BLOOM APPLICATION

Pests	<i>Belay</i> Insecticide Application Rate	Special Instructions
Blackheaded Fireworm (suppression) Cranberry Fruitworm (suppression) Cranberry Weevil Flea Beetles Leafhoppers Sparganothis Fruitworm (suppression) Stinkbugs Tipworm (suppression)	4 fl oz/A (0.067 lb ai/A)	Apply <i>Belay</i> Insecticide when target pest(s) threshold populations are observed. Do not wait until population beyond threshold has been established Apply <i>Belay</i> Insecticide in sufficient water to ensure uniform and thorough coverage of foliage. Thorough coverage is required for control.

- Do not apply more than 12 fl oz (0.2 lb ai) of *Belay* Insecticide per acre per season.
- Do not incorporate an adjuvant with this product.
- Do not apply treatments less than 7 days apart.
- Do not apply within 21 days of harvest.
- *Belay* Insecticide must not be applied during bloom or if bees are actively foraging.

SOIL POST BLOOM APPLICATION

Pests	<i>Belay</i> Insecticide Application Rates	Special Instructions
Cranberry Girdler ¹ Cranberry Weevil Flea Beetles Tipworm White Grubs	12 fl oz/A (0.2 lb ai/A)	Make one post bloom soil application as indicated below. Irrigate with a minimum of 0.2 inch of water within 2 hours of application (if rainfall occurs within the 2 hours irrigation may not be necessary). Apply specified dose to soil by one of the following methods: 1. As a ground application soil spray directed to the root and crown area. 2. As a chemigation application.

- Do not apply more than 12 fl oz (0.2 lb ai) of *Belay* Insecticide per acre per season.
- Do not apply within 21 days of harvest.

¹Except California

POTATO AND SWEET POTATO		
IN-FURROW OR SIDE DRESS APPLICATIONS		
Pests	<i>Belay</i> Insecticide Application Rates	Special Instructions
Aphids Colorado Potato Beetles ¹ Corn Wireworm Flea Beetles Leafhoppers Potato Psyllids (suppression) Southern Potato Wireworm Sugarcane Beetle Sweet Potato Weevil Tobacco Wireworm White Grubs (including Exotic White Grub) Wireworms (except for Corn, Southern Potato and Tobacco Wireworm) (suppression)	9 to 12 fl oz/A (0.15 to 0.2 lb ai/A)	<p>Timing of Application: At planting or as a side-dress on both sides of the row at ground-crack during hilling. With ground-crack/hilling application make certain product is covered with at least 3 inches of soil.</p> <p>Apply specified rate in sufficient carrier volume to insure uniform application and incorporate into the soil using one of the following methods:</p> <ol style="list-style-type: none"> 1. In a narrow band centered on the plant row in the bedding operation just prior to planting. 2. In-furrow spray at planting. Direct spray in the furrow on the seed pieces or potatoes. 3. As a sidedress to both sides of the row or as a spray at ground crack directly over the row during hilling. Cover immediately with soil.
Diseases Suppressed	<i>Belay</i> Insecticide Application Rate	Chemigation from plant emergence to 50 percent row cover. Applications via chemigation targeting above ground pests after 50 percent row cover could result in limited plant uptake via the root system. Use at least 0.15 inches of water when chemigating <i>Belay</i> Insecticide.
Symptoms of: Potato Leafroll Virus (PLRV) Purple Top	12 fl oz/A (0.2 lb ai/A)	
<ul style="list-style-type: none"> • Regardless of formulation or type of application (seed piece, soil or foliar), do not apply more than a total of 0.2 lbs active ingredient clothianidin per acre per season. 		

¹*Belay* Insecticide will not control Colorado potato beetles in regions where insensitivity to neonicotinoid insecticides has been reported.

PLEASE CONTACT VALENT U.S.A. CORPORATION AT 800-6-VALENT (682-5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE.

Copyright© 2011 by Valent U.S.A. Corporation.

Products That Work, From People Who Care is a registered trademark of Valent U.S.A. Corporation. *Belay* is a registered trademark of Sumitomo Chemical Company, Limited

Manufactured for:

Valent U.S.A. Corporation / P.O. Box 8025 / Walnut Creek CA 94596-8025

Made in U.S.A.

Form 2011-BEL-0011

Products That Work, From People Who Care® | www.valent.com | 800-6-VALENT (682-5368)