

# Supplemental Label



## **BELAY<sup>®</sup>** **INSECTICIDE**

EPA Reg. No. 59639-150  
(Florida Only)

### **BELAY<sup>®</sup> INSECTICIDE FOR POTATO AND SWEET POTATO**

This supplemental label expires on October 2, 2012 and must not be used or distributed after this date.

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. READ THE LABEL AFFIXED TO THE CONTAINER FOR *BELAY* INSECTICIDE BEFORE APPLYING. USE OF *BELAY* INSECTICIDE ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR *BELAY* INSECTICIDE.**

| POTATO AND SWEET POTATO  |  |   |
|--|--|---|
| IN-FURROW OR SIDE DRESS APPLICATIONS   |  |   |
| Pests  | <i>Belay</i> Insecticide Application Rates       | Special Instructions  |
| Aphids<br>Colorado Potato Beetles <sup>1</sup><br>Corn Wireworm<br>Flea Beetles<br>Leafhoppers<br>Potato Psyllids (suppression)<br>Southern Potato Wireworm<br>Sugarcane Beetle<br>Sweet Potato Weevil<br>Tobacco Wireworm<br>White Grubs (including Exotic White Grub)<br>Wireworms (except for Corn, Southern Potato and Tobacco Wireworm) (suppression) | 9 to 12 fl oz/A<br>(0.15 to 0.2 lb ai/A)         | <p><b>Timing of Application:</b> At planting or as a side-dress on both sides of the row at ground-crack during hilling. With ground-crack/hilling application make certain product is covered with at least 3 inches of soil.</p> <p>Apply specified rate in sufficient carrier volume to insure uniform application and incorporate into the soil using one of the following methods:</p> <ol style="list-style-type: none"> <li>1. In a narrow band centered on the plant row in the bedding operation just prior to planting.</li> <li>2. In-furrow spray at planting. Direct spray in the furrow on the seed pieces or potatoes.</li> <li>3. As a sidedress to both sides of the row or as a spray at ground crack directly over the row during hilling. Cover immediately with soil.</li> </ol> |
| <b>Diseases Suppressed</b>   | <b><i>Belay</i> Insecticide Application Rate</b> | Chemigation from plant emergence to 50 percent row cover. Applications via chemigation targeting above ground pests after 50 percent row cover could result in limited plant uptake via the root system. Use at least 0.15 inches of water when chemigating <i>Belay</i> Insecticide.   |
| Symptoms of:<br>Potato Leafroll Virus (PLRV)<br>Purple Top   | 12 fl oz/A<br>(0.2 lb ai/A)                      |   |
| <ul style="list-style-type: none"> <li>• Regardless of formulation or type of application (seed piece, soil or foliar), do not apply more than a total of 0.2 lbs active ingredient clothianidin per acre per season.</li> </ul>   |  |   |

<sup>1</sup>*Belay* Insecticide will not control Colorado potato beetles in regions where insensitivity to neonicotinoid insecticides has been reported.

**PLEASE CONTACT VALENT U.S.A. CORPORATION AT 800-6-VALENT (682-5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE.**

Copyright© 2011 by Valent U.S.A. Corporation.

*Belay* is a registered trademark of Sumitomo Chemical Company, Limited

Manufactured for:  
**Valent U.S.A. Corporation**  
P.O. Box 8025  
Walnut Creek CA 94596 - 8025  
(800) 682-5368  
Made in U.S.A.