

# Supplemental Label



## **BELAY<sup>®</sup>** **INSECTICIDE**

EPA Reg. No. 59639-150  
(Florida Only)

**BELAY<sup>®</sup> INSECTICIDE FOR CONTROL OF LISTED SUCKING AND CHEWING INSECTS INFESTING COTTON, CRANBERRY (LOW-GROWING BERRY, EXCEPT STRAWBERRY), FIG, GRAPE, PEACH, POME FRUIT, POMEGRANATE, SOYBEAN, TOBACCO, TREE NUTS AND TUBEROUS AND CORM VEGETABLES (INCLUDING POTATO AND SWEET POTATO).**

This supplemental label expires on October 2, 2012 and must not be used or distributed after this date.

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. READ THE LABEL AFFIXED TO THE CONTAINER FOR *BELAY* INSECTICIDE BEFORE APPLYING. USE OF *BELAY* INSECTICIDE ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR *BELAY* INSECTICIDE.**

### **RESTRICTIONS**

*BELAY* Insecticide is an insecticide for broad spectrum control of insects infesting cotton, cranberry (low-growing berry, except strawberry, fig, grape, peach, pome fruit, pomegranate, soybean, tobacco, tree nuts and tuberous and corm vegetables (including potato and sweet potato). When applied as directed, *BELAY* Insecticide provides excellent and long residual control.

- Regardless of formulation or type of application (seed treatment, soil or foliar), do not apply more than a total of 0.2 lbs active ingredient clothianidin per acre per season for all crops.
- If the maximum season limit of clothianidin as mentioned above has been applied and pest populations require additional treatments, use another registered pesticide that is not in the neonicotinoid class of chemistry.
- Do not apply by air except for cotton, potato and soybean.
- Do not incorporate an adjuvant with this product on almond, cranberry (low-growing berry, except strawberry), and peach.

ROTATIONAL CROP PLANT-BACK INTERVALS				
Crops Planted	Immediate Plant-Back	30-day Plant-Back	8 Month Plant-Back	12 Month Plant-Back
Cotton, Soybean and Tuberous and Corm Vegetables	Cereal Grains (except Rice), Cotton, Fruiting Vegetables (including Cucurbits), Leafy Vegetables (including Brassica Vegetables), Soybean, Bulb Vegetables, Root and Tuber Vegetables, Rapeseed, Sorghum, Canola, Corn (Field, Pop and Sweet)	Dry Beans, Grasses, Non-grass Animal Feeds, Rice	Sugarcane Any	crops without an earlier plant-back interval

COTTON		
FOLIAR APPLICATION		
Pests	<i>BELAY</i> Insecticide Application Rates	Special Instructions
Aphids <sup>1</sup> Flea Beetles Fleahoppers Lygus/Plant Bugs Stinkbugs Thrips (suppression) Whiteflies (suppression)	3 to 6 fl oz/A (0.05 to 0.1 lb ai/A)	Apply <i>BELAY</i> Insecticide when target pest(s) threshold populations are observed. Do not wait until population beyond threshold has been established.  Apply <i>BELAY</i> Insecticide in sufficient water to ensure uniform and thorough coverage of foliage. Thorough coverage is required for control. Use between 3 to 10 gallons per acre for aerial applications.
<ul style="list-style-type: none"> <li>Do not apply more than 12 fl oz (0.2 lb ai) of <i>BELAY</i> Insecticide per acre per season.</li> <li>Do not apply treatments less than 7 days apart.</li> <li>Do not apply within 21 days of harvest.</li> </ul>		

<sup>1</sup> Heavy infestations of aphids might require two applications of the highest rate to achieve control. *BELAY* Insecticide will not control cotton aphids in regions where insensitivity to neonicotinoid insecticides has been reported.

**CRANBERRY (LOW-GROWING BERRY EXCEPT STRAWBERRY)**

Bearberry, Bilberry, Cloudberry, Cranberry, Lingonberry, Lowbush Blueberry, Muntries, Partridgeberry and varieties, and/or cultivars of these

**FOLIAR POST BLOOM APPLICATION**

<b>Pests</b>	<b><i>BELAY</i> Insecticide Application Rate</b>	<b>Special Instructions</b>
Blackheaded Fireworm (suppression) Cranberry Fruitworm (suppression) Cranberry Weevil Flea Beetles Leafhoppers Sparganothis Fruitworm (suppression) Stinkbugs Tipworm (suppression)	4 fl oz/A (0.067 lb ai/A)	Apply <i>BELAY</i> Insecticide when target pest(s) threshold populations are observed. Do not wait until population beyond threshold has been established  Apply <i>BELAY</i> Insecticide in sufficient water to ensure uniform and thorough coverage of foliage. Thorough coverage is required for control.

- Do not apply more than 12 fl oz (0.2 lb ai) of *BELAY* Insecticide per acre per season.
- Do not incorporate an adjuvant with this product.
- Do not apply treatments less than 7 days apart.
- Do not apply within 21 days of harvest.
- *BELAY* Insecticide must not be applied during bloom or if bees are actively foraging.

**SOIL POST BLOOM APPLICATION**

<b>Pests</b>	<b><i>BELAY</i> Insecticide Application Rate</b>	<b>Special Instructions</b>
Cranberry Girdler <sup>1</sup> Cranberry Weevil Flea Beetles Tipworm White Grubs	12 fl oz/A (0.2 lb ai/A)	Make one post bloom soil application as indicated below. Irrigate with a minimum of 0.2 inch of water within 2 hours of application (if rainfall occurs within the 2 hours irrigation may not be necessary).  <b>Apply specified dose to soil by one of the following methods:</b> 1. As a ground application soil spray directed to the root and crown area. 2. As a chemigation application.

- Do not apply more than 12 fl oz (0.2 lb ai) of *BELAY* Insecticide per acre per season.
- Do not apply within 21 days of harvest.

<sup>1</sup>Except California

**FIG**

**FOLIAR POST BLOOM APPLICATION**

<b>Pests</b>	<b><i>BELAY</i> Insecticide Application Rates</b>	<b>Special Instructions</b>
Dried Fruit Beetles Fig Scale	4 to 6 fl oz/A (0.067 to 0.1 lb ai/A)	<p>Apply <i>BELAY</i> Insecticide when target pest(s) threshold populations are observed. Do not wait until population beyond threshold has been established.</p> <p>The amount of <i>BELAY</i> Insecticide per acre will depend on tree size and pest pressure.</p> <p>Choose lower rate for light infestation and/or small trees and the higher rate for heavy infestations and/or larger trees.</p> <p>Apply by ground 100 to 400 gallons per acre as a full coverage spray. Spray volume will depend on tree size and density of canopy. Thorough coverage is required for control. To achieve optimum coverage, use proper spray pressures, nozzles, nozzle spacing and water volume per acre.</p> <p>Repeat applications as needed to maintain control but do not apply more often than every 14 days. A single application may result in suppression only.</p>

- Do not apply more than 12 fl oz (0.2 lb ai) of *BELAY* Insecticide per season.
- Do not apply treatments less than 14 days apart.
- Do not apply within 7 days of harvest.
- Do not feed or allow livestock to graze on cover crops from treated orchards.

<b>GRAPE</b>		
<b>FOLIAR APPLICATION</b>		
<b>Pests</b>	<b><i>BELAY</i> Insecticide Application Rates</b>	<b>Special Instructions</b>
Leafhoppers Grape Leaf Skeletonizers Japanese Beetles Multicolored Asian Lady Beetle	2 to 4 fl oz/A (0.033 to 0.067 lb ai/A)	Apply <i>BELAY</i> Insecticide when target pest(s) threshold populations are observed. Do not wait until population beyond threshold has been established.
Sharpshooters	4 to 6 fl oz/A (0.067 to 0.1lb ai/A)	Apply <i>BELAY</i> Insecticide in sufficient water to ensure uniform and thorough coverage of foliage. Thorough coverage is required for control
European Grapevine Moth Grape Berry Moth Grape Mealybug Longtailed Mealybug Obscure Mealybug Vine Mealybug	6 fl oz/A (0.1 lb ai/A)	
<ul style="list-style-type: none"> <li>• Do not apply more than 12 fl oz (0.2 lb ai) of <i>BELAY</i> Insecticide per acre per season.</li> <li>• Do not apply treatments less than 14 days apart.</li> <li>• May be applied up to harvest (0 days PHI).</li> </ul>		

continued

**GRAPE** (continued)

<b>SOIL APPLICATION</b>		
<b>Pests</b>	<b><i>BELAY</i> Insecticide Application Rates</b>	<b>Special Instructions</b>
Grape Mealybug Grape Phylloxera spp. (suppression) Leafhoppers Longtailed Mealybug Obscure Mealybug Sharpshooters Vine Mealybug	6 to 12 fl oz/A (0.1 to 0.2 lb ai/A)	Apply <i>BELAY</i> Insecticide when target pest(s) threshold populations are observed. Do not wait until population beyond threshold has been established. Apply specified rate in sufficient carrier volume to ensure uniform application and incorporate into the soil using drip or trickle irrigation water.  Make one soil application at specified dose in sufficient carrier volume to insure uniform application using the following method: <ul style="list-style-type: none"> <li>• Chemigation into root zone through drip, trickle or any similar custom made systems that target the watering of the root zone of the plants.</li> </ul>
<ul style="list-style-type: none"> <li>• Do not apply more than 12 fl oz (0.2 lb ai) of <i>BELAY</i> Insecticide per acre per season.</li> <li>• Do not apply treatments less than 14 days apart.</li> <li>• Do not apply within 30 days of harvest.</li> </ul>		

<b>PEACH</b>		
<b>FOLIAR POST BLOOM APPLICATION</b>		
<b>Pests</b>	<b><i>BELAY</i> Insecticide Application Rates</b>	<b>Special Instructions</b>
Aphids Leafhoppers	3 to 6 fl oz/A (0.05 to 0.1 lb ai/A)	Apply <i>BELAY</i> Insecticide when target pest(s) threshold populations are observed. Do not wait until population beyond threshold has been established.  Choose lower rate for light infestation and/or small trees and the higher rate for heavy infestations and/or larger trees.
Lygus/Plant Bugs Plum Curculio Scales Stinkbugs	6 fl oz/A (0.1 lb ai/A)	Apply by ground 100 to 400 gallons per acre as a full coverage spray. Thorough coverage is required for control. To achieve optimum coverage, use proper spray pressures, nozzles, nozzle spacing and water volume per acre.  <b>Scales:</b> Time applications to the crawler stage, treat each successive generation. Two applications on a 10- to 14- day interval may be required to achieve control.
<ul style="list-style-type: none"> <li>• Do not apply more than 12 fl oz (0.2 lb ai) of <i>BELAY</i> Insecticide per acre per season.</li> <li>• Do not incorporate an adjuvant with this product.</li> <li>• Do not apply treatments less than 10 days apart.</li> <li>• Do not apply within 21 days of harvest.</li> <li>• Do not feed or allow livestock to graze on cover crops from treated orchards.</li> <li>• <i>BELAY</i> Insecticide must not be applied during bloom or if bees are actively foraging.</li> </ul>		

**POME FRUIT**

Apple, Crabapple, Loquat, Mayhaw, Pear, Oriental Pear, Quince

**FOLIAR POST BLOOM APPLICATION**

<b>Pests</b>	<b><i>BELAY</i> Insecticide Application Rates</b>	<b>Special Instructions</b>
Aphids Leafhoppers Plant Bugs Stinkbugs	4 to 6 fl oz/A (0.067 to 0.1 lb ai/A)	Apply <i>BELAY</i> Insecticide when target pest(s) threshold populations are observed. Do not wait until population beyond threshold has been established.
Apple Maggot Leafminers Mealybugs Plum Curculio	6 fl oz/A (0.1 lb ai/A)	The amount of <i>BELAY</i> Insecticide per acre will depend on tree size and pest pressure. Choose lower rate for light infestation and/or small trees and the higher rate for heavy infestation and/or larger trees.
Codling Moth* Lygus Oblique-banded Leafroller Oriental Fruit Moth Pear Psylla Scales Stinkbugs	6 to 12 fl oz/A (0.1 to 0.2 lb ai/A)	<p>Apply by ground 100 to 400 gallons per acre as a full coverage spray. Thorough coverage is required for control. To achieve optimum coverage, use proper spray pressures, nozzles, nozzle spacing and water volume per acre.</p> <p>Applications may also be made as a concentrate spray. Maintain the same amount of product per acre as a dilute application using the appropriate amount of concentrate spray volume. Repeat applications as needed to maintain control but do not apply more often than every 10 days. A single application may result in suppression only.</p> <p>For optimum control of Leafminers and Leafrollers, make the first application as soon as pollination is complete and bees are removed from the orchard.</p> <p><b>*For Codling Moth (First Generation):</b> For control of first generation codling moth in areas with light pressure and suppression of first generation codling moth in areas of heavy infestation.</p> <p>Suppression only of second or third generation codling moth in any area.</p> <p><b>Scales:</b> Time applications to the crawler stage, treat each successive generation. Two applications on a 10- to 14-day interval may be required to achieve control.</p>

- Do not apply more than 12 fl oz (0.2 lb ai) of *BELAY* Insecticide per acre per season.
- Do not apply within 7 days of harvest.
- Do not feed or allow livestock to graze on cover crops from treated orchards.
- *BELAY* Insecticide must not be applied during bloom or if bees are actively foraging.

**POMEGRANATE**

**FOLIAR POST BLOOM APPLICATION**

<b>Pests</b>	<b><i>BELAY</i> Insecticide Application Rates</b>	<b>Special Instructions</b>
<p>Aphids Leafhoppers Mealybugs Whiteflies (suppression)</p>	<p>4 to 6 fl oz/A (0.067 to 0.1 lb ai/A)</p>	<p>Apply <i>BELAY</i> Insecticide when target pest(s) threshold populations are observed. Do not wait until population beyond threshold has been established.</p> <p>The amount of <i>BELAY</i> Insecticide per acre will depend on tree size and pest pressure.</p> <p>Choose lower rate for light infestation and/or small trees and the higher rate for heavy infestations and/or larger trees.</p> <p>Apply by ground 100 to 400 gallons per acre as a full coverage spray. Thorough coverage is required for control. To achieve optimum coverage, use proper spray pressures, nozzles, nozzle spacing and water volume per acre.</p> <p>Repeat applications as needed to maintain control but do not apply more often than every 14 days. A single application may result in suppression only.</p>
<ul style="list-style-type: none"> <li>• Do not apply more than 12 fl oz (0.2 lb ai) of <i>BELAY</i> Insecticide per season.</li> <li>• Do not apply treatments less than 14 days apart.</li> <li>• Do not apply within 7 days of harvest.</li> <li>• Do not feed or allow livestock to graze on cover crops from treated orchards.</li> <li>• <i>BELAY</i> Insecticide must not be applied during bloom or if bees are actively foraging.</li> </ul>		

<b>SOYBEAN</b>		
<b>FOLIAR APPLICATION</b>		
<b>Pests</b>	<b><i>BELAY</i> Insecticide Application Rates</b>	<b>Special Instructions</b>
Aphids <sup>1</sup> Bean Leaf Beetles Blister Beetles Japanese Beetles Leafhoppers Lygus/Plant Bugs Stinkbugs Three Cornered Alfalfa Hopper Whiteflies (suppression)	3 to 6 fl oz/A (0.05 to 0.1 lb ai/A)	Apply <i>BELAY</i> Insecticide when target pest(s) threshold populations are observed. Do not wait until population beyond threshold has been established.  Apply <i>BELAY</i> Insecticide in sufficient water to ensure uniform and thorough coverage of foliage. Thorough coverage is required for control. Use between 3 to 10 gallons per acre for aerial applications.
<ul style="list-style-type: none"> <li>• Do not apply foliar treatments less than 7 days apart.</li> <li>• Do not apply within 21 days of harvest.</li> <li>• Do not make foliar applications of <i>BELAY</i> Insecticide in fields treated with a neonicotinoid insecticide seed treatment(s) within 45 days after planting.</li> <li>• Regardless of formulation or type of application (foliar or seed treatment), do not apply more than 0.2 lb ai of clothianidin per acre per season.</li> <li>• Do not graze or feed soybean forage and hay to livestock.</li> </ul>		

<sup>1</sup> Applications for soybean aphid need to be targeted at or less than 250 aphids/plant. Populations over threshold might require more than one application to achieve control.

## TOBACCO

### FOLIAR APPLICATION

Pests	<i>BELAY</i> Insecticide Application Rates	Special Instructions
Aphids Flea Beetles	3 to 4 fl oz/A (0.05 to 0.067 lb ai/A)	Apply <i>BELAY</i> Insecticide when target pest(s) threshold populations are observed. Do not wait until population beyond threshold has been established.
Tobacco Budworm (suppression) Tobacco Hornworm (suppression)	4 fl oz/A (0.067 lb ai/A)	<p>Make application using equipment that delivers a coarse droplet spray and low pressure to prevent off target drift. Check calibration periodically to ensure that your equipment is working properly. Thorough coverage is required for control.</p> <p>For suppression of budworms and hornworms, make the first application as soon as eggs hatch and repeat applications as necessary to maintain suppression.</p>

- Do not apply more than 12 fl oz (0.2 lb ai) *BELAY* Insecticide per acre per season.
- Do not apply treatments less than 7 days apart.
- Do not apply within 14 days of harvest.

### SOIL APPLICATION

(In Transplanted Water Application or Tray, or Flat Drench Application)

Pests	<i>BELAY</i> Insecticide Application Rates	Special Instructions
Flea Beetles	3 to 6 fl oz/A (0.05 to 0.1 lb ai/A)	<p><i>BELAY</i> Insecticide must be mixed with water prior to application. Keep the <i>BELAY</i> Insecticide solution agitated or mix regularly to prevent settling in the spray or transplant tank. Apply <i>BELAY</i> Insecticide at the specified dosage to the soil in transplant water or by drench to trays or flats prior to transplanting. Water the plants in flats or trays by overhead irrigation to wash <i>BELAY</i> Insecticide from foliage into the potting media. Failure to wash <i>BELAY</i> Insecticide from foliage may result in a reduction of control.</p> <p>Soil applications may sometimes cause yellowing on the leaves present at the time of transplanting. This effect does not have any impact on plant growth and yields.</p>
Aphids Tobacco Budworm (suppression) Tobacco Hornworm (suppression)	6 fl oz./A (0.1 lb ai/A)	

- Do not apply more than 12 fl oz (0.2 lb ai) of *BELAY* Insecticide per acre per season
- Do not apply within 14 days of harvest.

### TREE NUTS

Almond, Beech Nut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (Hazelnut), Hickory Nut, Macadamia Nut, Pecan, Walnut (Black and English)

#### FOLIAR POST BLOOM APPLICATION

Pests	<i>BELAY</i> Insecticide Application Rates	Special Instructions
Aphids Leafhoppers Pecan Nut Casebearer Pecan Weevil Walnut Husk Fly Whiteflies (suppression)	3 to 6 fl oz/A (0.05 to 0.1 lb ai/A)	Apply <i>BELAY</i> Insecticide when target pest(s) threshold populations are observed. Do not wait until population beyond threshold has been established.  Choose lower rate for light infestation and/or small trees and the higher rate for heavy infestation and/or larger trees.
Hickory Shuckworm Lygus Mealybugs Scales Stinkbugs	6 fl oz/A (0.1 lb ai/A)	Apply by ground 100 to 400 gallons per acre as a full coverage spray. Thorough coverage is required for control. To achieve optimum coverage, use proper spray pressures, nozzles, nozzle spacing and water volume per acre.  <b>Scales:</b> Time applications to the crawler stage, treat each successive generation. Two applications on a 10 to 14-day interval may be required to achieve control.

- Do not apply more than 12 fl oz (0.2 lb ai) of *BELAY* Insecticide per acre per season.
- Do not incorporate an adjuvant with this product on almond.
- Do not apply treatments less than 10 days apart.
- Do not apply within 21 days of harvest.
- *BELAY* Insecticide must not be applied during bloom or if bees are actively foraging.
- Do not feed or allow livestock to graze on cover crops from treated orchards.

**TUBEROUS AND CORM VEGETABLES**

Arrachcha, arrowroot, artichoke, (including Chinese and Jerusalem), edible canna, cassava (including bitter and sweet), root chayote, chufa, dasheen, ginger, leren, potato, sweet potato, taniel, turmeric, and yam

**FOLIAR APPLICATION**

<b>Pests</b>	<b><i>BELAY</i> Insecticide Application Rates</b>	<b>Special Instructions</b>
Aphids Colorado Potato Beetles <sup>1</sup> Flea Beetles Leafhoppers	2 to 3 fl oz/A (0.033 to 0.05 lb ai/A)	Apply <i>BELAY</i> Insecticide when target pest(s) threshold populations are observed. Do not wait until population beyond threshold has been established. Thorough coverage is required for control. Use between 3 to 10 gallons per acre for aerial applications.

- Do not apply treatments less than 7 days apart.
- Do not apply within 14 days of harvest.
- Regardless of formulation or type of application (seed piece, soil or foliar), do not apply more than a total of 0.2 lbs active ingredient clothianidin per acre per season.
- Do not apply by air except for potato.

<sup>1</sup> *BELAY* Insecticide will not control Colorado potato beetles in regions where insensitivity to neonicotinoid insecticides has been reported.

continued

**TUBEROUS AND CORM VEGETABLES** (continued)

<b>IN-FURROW OR SIDE DRESS APPLICATIONS</b>		
<b>Pests</b>	<b><i>BELAY</i> Insecticide Application Rates</b>	<b>Special Instructions</b>
Aphids Colorado Potato Beetles <sup>1</sup> Corn Wireworm Flea Beetles Leafhoppers Potato Psyllids (suppression) Southern Potato Wireworm Sugarcane Beetle Sweet Potato Weevil Tobacco Wireworm White Grubs (including Exotic White Grub) Wireworms (except for Corn, Southern Potato and Tobacco Wireworm) (suppression)	9 to 12 fl oz/A (0.15 to 0.2 lb ai/A)	<p><b>Timing of Application:</b> At planting or as a side-dress on both sides of the row at ground-crack during hilling. With ground-crack/hilling application make certain product is covered with at least 3 inches of soil.</p> <p>Apply specified rate in sufficient carrier volume to insure uniform application and incorporate into the soil using one of the following methods:</p> <ol style="list-style-type: none"> <li>1. In a narrow band centered on the plant row in the bedding operation just prior to planting.</li> <li>2. In-furrow spray at planting. Direct spray in the furrow on the seed pieces or potatoes.</li> <li>3. As a sidedress to both sides of the row or as a spray at ground crack directly over the row during hilling. Cover immediately with soil.</li> </ol>
<b>Diseases Suppressed</b>	<b><i>BELAY</i> Insecticide Application Rate</b>	Chemigation from plant emergence to 50 percent row cover. Applications via chemigation targeting above ground pests after 50 percent row cover could result in limited plant uptake via the root system. Use at least 0.15 inches of water when chemigating <i>BELAY</i> Insecticide.
Symptoms of: Potato Leafroll Virus (PLRV) Purple Top	12 fl oz/A (0.2 lb ai/A)	
<ul style="list-style-type: none"> <li>• Regardless of formulation or type of application (seed piece, soil or foliar), do not apply more than a total of 0.2 lbs active ingredient clothianidin per acre per season.</li> </ul>		

<sup>1</sup> *BELAY* Insecticide will not control Colorado potato beetles in regions where insensitivity to neonicotinoid insecticides has been reported.

continued

**TUBEROUS AND CORM VEGETABLES (continued)**

<b>SEED-PIECE APPLICATION</b>		
<b>Pests</b>	<b><i>BELAY</i> Insecticide Rates/100 lbs seed</b>	<b>Special Instructions</b>
Aphids Colorado Potato Beetles <sup>1</sup> Flea Beetles Leafhoppers Potato Leafhoppers Potato Psyllids (suppression only) Wireworms (seed-piece protection only)	0.4 to 0.6 fl oz*	<b>Seed-Piece Application:</b> 1. Apply specified dosage as a diluted, fine spray applied over the cut or whole seed tubers. A dilution rate of 1 part <i>BELAY</i> Insecticide to 3 parts water, or less, is recommended. 2. Maintain agitation of spray mixture during the application process. 3. Use a spray system that is properly shielded to prevent any spray from moving off target.
<b>Diseases Suppressed</b>	<b><i>BELAY</i> Insecticide Rates/100 lbs seed</b>	
Symptoms of: Potato Leaf Roll virus (PLRV) Purple Top Net Necrosis	0.6 fl oz	Liquid-based fungicides can be applied with <i>BELAY</i> Insecticide; however check compatibility before use. If using inert dusts or dust-based fungicides, apply <i>BELAY</i> Insecticide first, followed by dust application.  Apply only in areas with adequate ventilation to accommodate dry-down and in areas equipped to remove mist or dust.  Plant treated seed-pieces as soon as possible after treating.  *Based on seeding rate of 2,000 lbs/acre, seed-piece application rate range equals in-furrow/side-dress per acre application rate range.
<ul style="list-style-type: none"> <li>• Do not use treated seed-pieces for food, feed or fodder.</li> <li>• Do not apply any subsequent application of <i>BELAY</i> Insecticide (in-furrow, side-dress) following a <i>BELAY</i> Insecticide seed-piece treatment.</li> <li>• Regardless of formulation or type of application (seed piece, soil or foliar), do not apply more than a total of 0.2 lbs active ingredient clothianidin per acre per season.</li> </ul>		

<sup>1</sup> *BELAY* Insecticide will not control Colorado Potato Beetles in regions where insensitivity to neonicotinoid insecticides has been reported.

**PLEASE CONTACT VALENT U.S.A. CORPORATION AT 800-6-VALENT (682-5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE.**

Copyright© 2011 by Valent U.S.A. Corporation.

*Products That Work, From People Who Care* is a registered trademark of Valent U.S.A. Corporation.  
*Belay* is a registered trademark of Sumitomo Chemical Company, Limited

Manufactured for:

**Valent U.S.A. Corporation**

P.O. Box 8025

Walnut Creek CA 94596 - 8025

Made in U.S.A.

Form 2011-BEL-0013

20110310

*Products That Work, From People Who Care*® | [www.valent.com](http://www.valent.com) | 800-6-VALENT (682-5368)