

# Supplemental Label



EPA Reg. No. 59639-119  
(Except New York)

## CHATEAU<sup>®</sup> HERBICIDE WDG USE IN Highbush Blueberry

THIS SUPPLEMENTAL LABEL SUPERCEDES PREVIOUS LABELING FOR USE IN Highbush Blueberry.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. READ THE LABEL AFFIXED TO THE CONTAINER FOR *CHATEAU* HERBICIDE WDG BEFORE APPLYING. USE OF *CHATEAU* HERBICIDE WDG ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR *CHATEAU* HERBICIDE WDG.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.**

### USE PRECAUTIONS

- Do not use in the states of Idaho, Oregon or Washington except west of the Cascade Mountains in the following counties:  
**Oregon:** Benton, Clackamas, Clatsop, Columbia, Coos, Curry, Douglas, Jackson, Josephine, Lane, Lincoln, Linn, Marion, Multnomah, Polk, Tillamook, Yamhill and Washington.  
**Washington:** Clallam, Cowlitz, Grays Harbor, King, Jefferson, Kitsap, Lewis, Pacific, Pierce, Skagit, Snohomish, Thurston, Wahkiakum, Whatcom..
- Do not apply to bushberries established less than 2 years unless they are protected from spray contact by non-porous wrap, grow tubes or waxed containers.
- Do not apply within 7 days of harvest.

### RESTRICTIONS AND LIMITATIONS

- Do not apply more than 12 oz of *Chateau* Herbicide WDG per acre during a single application.
- Do not apply more than 12 oz of *Chateau* Herbicide WDG per acre during a 12 month period.
- Do not make a sequential application within 30 days of the first application.
- A maximum *Chateau Herbicide WDG* rate of 6 oz/A per application should be used on any soil that has a sand plus gravel content over 80% if bushes are less than 3 years of age. (Two applications of 6 oz/A in a 12 month period can still be made as long as there have been 60 days between applications).
- Do not apply to farm alleys or roads where traffic may result in treated dust settling onto crops or other desirable vegetation.
- Do not apply within 300 yards of non-dormant pears.
- Raise mower height during all mowing to reduce dust. Dust created by mowing can drift onto desirable vegetation resulting in injury.

- Do not apply to powdery soils or soils that are susceptible to wind displacement unless irrigation can be applied immediately after application.
- Do not mow treated areas between bud break and final harvest. Dust created by mowing may drift onto desirable vegetation resulting in injury.
- Follow the most restrictive label limitations and precautions of the tank mix product(s) being used.
- Avoid direct or indirect spray contact to foliage and green bark (non-barked trunk with the exception of undesirable suckers).

*Chateau Herbicide WDG should be applied as a uniform band directed at the base of the bush. The preferred application timing for Chateau Herbicide WDG is in the fall to maximize the potential for rainfall to activate and set the herbicide. Do not apply over the top of crop or allow spray to come in contact with crop as a result of application or drift.*

### **Preemergence Application**

Apply 6 to 12 oz (0.188 to 0.38 lb ai/A) of *Chateau Herbicide WDG* per broadcast acre as a pre-emergence application. Preemergence (to weed emergence) applications of *Chateau Herbicide WDG should be made to a weed-free soil surface. Preemergence applications of Chateau Herbicide WDG must be completed prior to weed emergence. Moisture is necessary to activate Chateau Herbicide WDG on soil for residual weed control. Dry weather following application of Chateau Herbicide WDG may reduce effectiveness. However, when adequate moisture is received after dry conditions, Chateau Herbicide WDG will control susceptible germinating weeds.*

### **Postemergence Application**

Apply 6 to 12 oz (0.188 to 0.38 lb ai/A) of *Chateau Herbicide WDG* per broadcast acre plus an adjuvant (0.25% v/v non-ionic surfactant or 1 qt./A crop oil concentrate). The addition of an adjuvant enhances *Chateau Herbicide WDG* activity on emerged weeds. Thorough spray coverage is necessary to maximize the postemergence activity of *Chateau Herbicide WDG*.

Refer to Table 1 for weeds controlled by the residual activity of *Chateau Herbicide WDG*. *Chateau Herbicide WDG should be tank mixed with a labeled burndown herbicide for control of the emerged weeds listed in Table 2. Refer to tank mix partner's label for additional weed species and increased weed heights claimed. Refer to tank mix partner's label for additional restrictions, including minimum carrier volume and crops in which tank mix partner may be used. Burndown tank mix partners include glyphosate, paraquat, 2,4-D and RELY®. Tank mixes with glyphosate or 2,4-D containing products are not recommended during the period after bloom through final harvest to ensure crop safety from drift.*

Residual weed control will be reduced if vegetation prevents the *Chateau Herbicide WDG* from reaching the soil surface. If vegetation is heavy, it is recommended to use a burndown herbicide with *Chateau Herbicide WDG* and make a sequential *Chateau Herbicide WDG* application prior to the emergence of new weeds.

### **Carrier Volume and Spray Pressure**

To ensure thorough coverage in burndown applications, use a minimum of 15 gallons of spray solution per acre. Use higher gallonage if dense vegetation or heavy crop residue is present.

Nozzle selection should meet manufacturer's gallonage and pressure recommendations.

### Banded Application

Rates listed in Table 2, Weeds Controlled by Postemergence Activity of *Chateau* Herbicide WDG Tank Mixes, refer to a broadcast application covering the entire acre. When making a banded application, the rate must be reduced according to the following formula:

Amount Needed per Acre for Banded Application	=	$\frac{\text{Band Width in inches}}{\text{Row Width in inches}}$	X	Rate per Broadcast Acre
---	---	--	---	-------------------------

**Table 1. Weeds Controlled by Preemergence Application of *Chateau* Herbicide WDG**

<b>BROADLEAF WEED SPECIES</b>				
COMMON NAME	SCIENTIFIC NAME	ORGANIC MATTER	SOIL TYPE	CHATEAU HERBICIDE WDG RATE
Bristly Starbur	<i>Acanthospermum hispidum</i>	Up to 10% <sup>1</sup>	All Soil Types <sup>2</sup>	6 to 12 oz/A <sup>2</sup>
Carpetweed	<i>Mollugo verticillata</i>			
Chickweeds				
Common	<i>Stellaria media</i>			
Mouseear	<i>Cerastium vulgatum</i>			
Coffee Senna	<i>Cassia occidentalis</i>			
Dandelion	<i>Taraxacum officinale</i>			
Eclipta	<i>Eclipta prostrata</i>			
Eveningprimrose, Cutleaf	<i>Oenothera laciniata</i>			
Filaree				
Redstem	<i>Erodium cicutarium</i>			
Whitestem	<i>Erodium moschatum</i>			
Fleabane	<i>Erigeron</i> spp.			
Florida Beggarweed	<i>Desmodium tortuosum</i>			
Florida Pusley	<i>Richardia scabra</i>			
Golden Crownbeard	<i>Verbesina encelioides</i>			
Groundsel, Common	<i>Senecio vulgaris</i>			
Hairy Indigo	<i>Indigofera hirsute</i>			
Hemp Sesbania	<i>Sesbania exaltata</i>			
Henbit	<i>Lamium amplexicaule</i>			
Jimsonweed	<i>Datura stramonium</i>			
Kochia	<i>Kochia scoparia</i>			
Lambsquarters, Common	<i>Chenopodium album</i>			
Mallow				
Common (Cheeseweed)	<i>Malva neglecta</i>			
Little Mallow	<i>Malva parviflora</i>			
Horseweed/Marestail	<i>Coryza Canadensis</i>			

continued

<sup>1</sup>*Chateau Herbicide WDG* can be used on soils with greater than 10% organic matter; however, length of residual control may be shorter than on soils with lower organic matter content.

<sup>2</sup>A maximum *Chateau Herbicide WDG* rate of 6 oz/A per application should be used on any soil that has a sand plus gravel content over 80% if bushes are under 3 years of age.

**Table 1. Weeds Controlled by Preemergence Application of *Chateau Herbicide WDG* (continued)**

<b>BROADLEAF WEED SPECIES</b>				
<b>COMMON NAME</b>	<b>SCIENTIFIC NAME</b>	<b>ORGANIC MATTER</b>	<b>SOIL TYPE</b>	<b>CHATEAU HERBICIDE WDG RATE</b>
Morningglories		Up to 10% <sup>1</sup>	All Soil Types <sup>2</sup>	6 to 12 oz/A <sup>2</sup>
Entireleaf	<i>Ipomoea hederacea</i> var. <i>integriuscula</i>			
Ivyleaf	<i>Ipomoea hederacea</i>			
Red/Scarlet	<i>Ipomoea coccinea</i>			
Smallflower	<i>Jacquemontia tamnifolia</i>			
Tall	<i>Ipomoea purpurea</i>			
Mustards				
Tumble	<i>Sisymbrium altissimum</i>			
Wild	<i>Brassica kaber</i>			
Nightshades				
Black	<i>Solanum nigrum</i>			
Eastern Black	<i>Solanum ptycanthum</i>			
Hairy	<i>Solanum sarrachoides</i>			
Pigweeds				
Palmer Amaranth	<i>Amaranthus palmeri</i>			
Redroot	<i>Amaranthus retroflexus</i>			
Smooth	<i>Amaranthus hybridus</i>			
Spiny Amaranth	<i>Amaranthus spinosus</i>			
Tumble	<i>Amaranthus albus</i>			
Prickly Lettuce (China Lettuce)	<i>Lactuca serriola</i>			
Prickly Sida (Teaweed)	<i>Sida spinosa</i>			
Puncturevine	<i>Tribulus terrestris</i>			
Purslane, Common	<i>Portulaca oleracea</i>			
Radish, Wild	<i>Raphanus raphanistrum</i>			
Ragweed, Common	<i>Ambrosia artemisiifolia</i>			
Redmaids	<i>Calandrinia ciliata</i> var. <i>menziessi</i> .			
Redweed	<i>Melochia corchorifolia</i>			
Shepherd's-purse	<i>Capsella bursa-pastoris</i>			
Sowthistle, Annual	<i>Sonchus oleraceus</i>			
Spotted Spurge	<i>Euphorbia maculata</i>			
Spurred Anoda	<i>Anoda cristata</i>			
Thistle, Russian	<i>Salsola iberica</i>			
Tropic Croton	<i>Croton glandulosus</i>			
Venice Mallow	<i>Hibiscus trionum</i>			

continued

<sup>1</sup> *Chateau Herbicide WDG* can be used on soils with greater than 10% organic matter; however, length of residual control may be shorter than on soils with lower organic matter content.

<sup>2</sup> A maximum *Chateau Herbicide WDG* rate of 6 oz/A per application should be used on any soil that has a sand plus gravel content over 80% if bushes are under 3 years of age.

**Table 1. Weeds Controlled by Preemergence Application of *Chateau Herbicide WDG* (continued)**

<b>BROADLEAF WEED SPECIES</b>				
<b>COMMON NAME</b>	<b>SCIENTIFIC NAME</b>	<b>ORGANIC MATTER</b>	<b>SOIL TYPE</b>	<b>CHATEAU HERBICIDE WDG RATE</b>
Waterhemp		Up to 10% <sup>1</sup>	All Soil Types <sup>2</sup>	6 to 12 oz/A <sup>2</sup>
Common	<i>Amaranthus rudis</i>			
Tall	<i>Amaranthus tuberculatus</i>			
Wild Poinsettia	<i>Euphorbia heterophylla</i>			
Wormwood, Biennial	<i>Artemisia biennis</i>			
<b>GRASS WEED SPECIES</b>				
Barnyardgrass	<i>Echinochloa crus-galli</i>			
Bluegrass, Annual	<i>Poa annua</i>			
California Lovegrass	<i>Eragrostis diffusa</i>			
Crabgrass				
Large	<i>Digitaria sanguinalis</i>			
Smooth	<i>Digitaria ischaemum</i>			
Foxtails				
Bristly	<i>Setaria verticillata</i>			
Giant	<i>Setaria faberi</i>			
Green	<i>Setaria viridis</i>			
Yellow	<i>Setaria glauca</i>			
Goosegrass	<i>Eleusine indica</i>			
Guineagrass	<i>Panicum maximum</i>			
Johnsongrass, Seedling	<i>Sorghum halepense</i>			
Signalgrass, Broadleaf	<i>Brachiaria platyphylla</i>			
Panicum				
Fall	<i>Panicum dichotomiflorum</i>			
Texas	<i>Panicum texaum</i>			

<sup>1</sup>*Chateau Herbicide WDG* can be used on soils with greater than 10% organic matter; however, length of residual control may be shorter than on soils with lower organic matter content.

<sup>2</sup>A maximum *Chateau Herbicide WDG* rate of 6 oz/A per application should be used on any soil that has a sand plus gravel content over 80% if bushes are under 3 years of age.

**Table 2. Weeds Controlled by Postemergence Activity of CHATEAU HERBICIDE WDG Tank Mixes**

<b>BROADLEAF WEED SPECIES</b>			
<b>COMMON NAME</b>	<b>SCIENTIFIC NAME</b>	<b>WEED HEIGHT/LENGTH (inches)</b>	<b>CHATEAU HERBICIDE WDG RATE</b>
Bindweed, Field <sup>1</sup>	<i>Convolvulus arvensis</i>	8	6 to 12 oz/A
Carpetweed	<i>Mollugo verticillata</i>	4	
Chickweeds			
Common	<i>Stellaria media</i>	4	
Mouseear	<i>Cerastium vulgatum</i>	4	
Cocklebur, Common	<i>Xanthium strumarium</i>	4	
Eveningprimrose, Cutleaf <sup>2</sup>	<i>Oenothera laciniata</i>	12	
Filaree			
Broadleaf	<i>Erodium botrys</i>	4	
Redstem	<i>Erodium cicutarium</i>	4	
Florida Beggarweed	<i>Desmodium tortuosum</i>	2	
Hemp Sesbania	<i>Sesbania exaltata</i>	8	
Jimsonweed	<i>Datura stramonium</i>	4	
Lambsquarters, Common	<i>Chenopodium album</i>	4	
Morningglories			
Entireleaf	<i>Ipomoea hederacea</i> var. <i>integriuscula</i>	4	
Ivyleaf	<i>Ipomoea hederacea</i>	4	
Pitted	<i>Ipomoea lacunosa</i>	6	
Red/Scarlet	<i>Ipomoea coccinea</i>	4	
Tall	<i>Ipomoea purpurea</i>	4	
Pigweeds			
Palmer Amaranth	<i>Amaranthus palmeri</i>	6	
Redroot	<i>Amaranthus retroflexus</i>	6	
Smooth	<i>Amaranthus hybridus</i>	6	
Plaintain, Broadleaf	<i>Plantago major</i>	6	
Prickly Sida (Teaweed)	<i>Sida spinosa</i>	6	
Purslanes			
Common	<i>Portulaca oleracea</i>	4	
Rock	<i>Calandrinia</i> spp.	2	
Ragweeds			
Common	<i>Ambrosia artemisiifolia</i>	2	
Giant	<i>Ambrosia trifida</i>	4	
Sicklepod	<i>Senna obtusifolia</i>	4	

continued

<sup>1</sup> Chateau Herbicide WDG will only provide control of the above ground portion of bindweed. Repeated applications will be needed to control regrowth.

<sup>2</sup> For acceptable control, cutleaf evening primrose should be 12 inches or less and in the rosette stage. Crop oil concentrate, at 1 pt/A, or non-ionic surfactant at 0.25% v/v, should be added to glyphosate tank mixes for cutleaf evening primrose control, including glyphosate formulations that contain a built-in adjuvant system.

**Table 2. Weeds Controlled by Postemergence Activity of *Chateau Herbicide WDG Tank Mixes***  
(continued)

<b>BROADLEAF WEED SPECIES</b>			
<b>COMMON NAME</b>	<b>SCIENTIFIC NAME</b>	<b>WEED HEIGHT/LENGTH (inches)</b>	<b>CHATEAU HERBICIDE WDG RATE</b>
Smartweeds			6 to 12 oz/A
Ladysthumb	<i>Polygonum persicaria</i>	4	
Pale	<i>Polygonum lapathifolium</i>	4	
Pennsylvania	<i>Polygonum pennsylvanicum</i>	4	
Rice Flatsedge	<i>Cyperus iria</i>	4	
Spotted Spurge	<i>Euphorbia maculata</i>	4	
Velvetleaf	<i>Abutilon theophrasti</i>	4	
Venice Mallow	<i>Hibiscus trionum</i>	4	
Waterhemp			
Common	<i>Amaranthus rudis</i>	2	
Tall	<i>Amaranthus tuberculatus</i>	2	
Wild Mustard	<i>Brassica kaber</i>	6	

**PLEASE CONTACT VALENT U.S.A. CORPORATION AT 800-6-VALENT (682-5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE.**

Copyright © 2011 by Valent U.S.A. Corporation

*Chateau* and Products That Work, From People Who Care are registered trademarks of Valent U.S.A. Corporation

Manufactured for:  
Valent U.S.A. Corporation  
P. O. Box 8025  
Walnut Creek, CA 94596-8025

Made in U.S.A.

Form 2011-CHA-0015

*Products That Work, From People Who Care*® | **kkk j UYbHWa** | 800-6-VALENT (682-5368)