



Supplemental Label

Domark[®] 230 ME Fungicide

EPA Reg. No. 80289-7

FOR USE ON FIELD CORN, POPCORN, CORN GROWN FOR SEED PRODUCTION

(Only for use in the following states: Alabama, Arkansas, Delaware, Florida, Georgia, Iowa, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maryland, Michigan, Minnesota, Missouri, Mississippi, North Carolina, North Dakota, Nebraska, New Jersey, Ohio, Oklahoma, Pennsylvania, South Carolina, South Dakota, Tennessee, Texas, Virginia, Wisconsin and West Virginia.)

This supplemental label expires December 31, 2014 and must not be used or distributed after this date.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. READ THE LABEL AFFIXED TO THE CONTAINER FOR *DOMARK* BEFORE APPLYING. USE OF *DOMARK* ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR *DOMARK*.

CROP USE RATES AND TIMING OF APPLICATIONS

Field Corn, Popcorn, Corn Grown For Seed Production

Disease	Dosage Rate		When to Apply	Special Use Instructions
	Fl oz/A	GPA		
Gray Leaf Spot (<i>Cercospora zeae-maydis</i>) Rust, Common (<i>Puccinia sorghi</i>) Rust, Southern (<i>Puccinia polysora</i>) Anthracnose Leaf Blight (<i>Colletotrichum graminicola</i>) Eye Spot (<i>Aureobasidium zeae</i>) Northern Corn Leaf Blight (<i>Exserohilum turcicum</i>) Northern Corn Leaf Spot (<i>Bipolaris zeicola</i>) Southern Corn Leaf Blight (<i>Bipolaris maydis</i>)	4.0 to 6.0 (0.060 to 0.090 lb ai/A)	Ground: Minimum of 10 GPA Aerial: Minimum of 2 GPA	Apply prior to disease onset when conditions favor disease development. Curative applications are most effective when disease incidence does not exceed 5% of the plants at time of application.	Use <i>Domark</i> as part of an Integrated Pest Management program (IPM). Apply as a foliar spray or via chemigation in sufficient water to obtain thorough coverage of plants. To limit the potential for resistance development, do not apply more than 6 fl oz per acre per year.

RESTRICTIONS AND LIMITATIONS

1. **Do not** make more than (1) application per year.
2. **Do not** apply more than 6 fl oz (0.090 lb ai tetraconazole) of *Domark* per acre per year.
3. **Do not** apply *Domark* after corn growth stage R-3 (milk).
4. **Do not** use adjuvants in sprays made between V-8 (8 leaf collar) and VT (lowest branch of the tassel visible but silks have not emerged) growth stage. A compatibility agent, another fungicide, or an insecticide may be included if needed and labeled for use in corn. Refer to adjuvant product label for specific use directions and restrictions. Always follow the more restrictive label.

PLEASE CONTACT VALENT U.S.A. CORPORATION AT 800-6-VALENT (682-5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE.

Copyright © 2011 by Valent U.S.A. Corporation

Domark is a registered trademark of Isagro S.p.A.

Manufactured by



Isagro USA, Inc.
430 Davis Drive, Suite 240
Morrisville, NC 27560

Manufactured for
Valent U.S.A. Corporation
P.O. Box 8025
Walnut Creek, CA 94596-8025

Form 2011-DOM-0010