

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF FLORIDA
COBRA® Herbicide
FOR CONTROL OF BROADLEAF IN PINE SEEDLINGS
EPA Reg. No. 59639-34
EPA SLN No. FL-890032

Active Ingredient	By Wt.
*Lactofen	23.2%
Inert Ingredients	76.8%
* 1-(carboethoxy)ethyl 5-[2-chloro-4-(trifluoromethyl) phenoxy]-2-nitrobenzoate	
Contains 2 lbs. active ingredient per gallon.	

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

Pine Seedlings Nurseries Only

SOUTHERN PINE SEEDBEDS

For control of various annual broadleaf weeds including Common Purslane, Eclipta, Florida Pusley, and Poorjoe.

Apply as a preemergence spray. Apply up to 1.6 pts. of COBRA® Herbicide per acre (0.4 lb. active ingredient per acre). Apply COBRA Herbicide within 48 hours after sowing. Use low pressure (30 to 50 psi) with 20 to 40 gals. of water after application. Do not apply more than 1.6 pts. of COBRA Herbicide during the sowing season.

* * * * *

Follow all applicable directions, restrictions and precautions on the EPA registered label.

This labeling must be in the possession of the user at the time of pesticide application.

24(c) Registrant:

VALENT U.S.A. Corporation
P.O. Box 8025
Walnut Creek, CA 94596-8025