

**FOR DISTRIBUTION AND USE ONLY IN THE STATE OF FLORIDA**  
**BOLERO7 8 EC**  
**FOR CONTROL OF JUNGLERICE, BARNYARDGRASS & GOOSEGRASS IN RICE**  
**EPA Reg. No. 59639-79**  
**Formerly EPA Reg. No. 63588-5-59639**  
**EPA SLN No. FL-930010**

Active Ingredient	By Wt.
*Thiobencarb .....	84%
Inert Ingredients .....	16%
*S-[(4-chlorophenyl)methyl] diethylcarbamothioate	
Contains 8 pounds S-[(4-chlorophenyl)methyl]	
diethylcarbamothioate per gal.	

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**Do not apply this product through any type of irrigation system.**

**RICE - FOR DRY SEEDED RICE**

**Late Preemergence** Application (Prior to rice and grass emergence) - For control of Jungle Rice and Barnyardgrass (Echinochloa spp.) and Goosegrass (Eleusine indica). Apply BOLERO7 8 EC 1 - 5 days prior to rice emergence; generally this would be 5 - 9 days after planting of the rice. Apply 4 pts. of BOLERO7 8 EC per acre after soils have been sealed by rainfall or by flushing. Apply to well prepared seedbeds free of large clods. Soil should be wet at time of application. Do not apply BOLERO7 8 EC as a preemergence treatment to cracked soil. Do not apply BOLERO7 8 EC to exposed rice seed.

**Early Postemergence** Application - For control of emerged Jungle Rice and Barnyardgrass. Apply 4 pts. of BOLERO7 8 EC per acre when Jungle Rice or Barnyardgrass has developed no further than the two-leaf stage. Rice may be fully emerged at time of application. Soil should be wet at time of application.

**Do Not Apply BOLERO7 8 EC to Second Crop (Stubble Crop) Rice**

## WATER MANAGEMENT

After application, flush the fields as often as necessary to prevent crusting and drying of the soil. Fields should be flooded as soon as the rice plants will tolerate permanent flooding.

## APPLICATION EQUIPMENT

**Aircraft:** Apply BOLERO7 8 EC in no less than 10 gals. spray mix per acre.

**Ground Sprayers:** Apply 20 - 50 gals. of total spray mix per acre.

## PRECAUTIONS

Do not plant subsequent crops in treated fields within 6 months of last application. In areas where double cropping is the agricultural practice, do not apply to a second stubble crop. Avoid drift to non-target areas.

Water drained directly from treated fields must not be used to irrigate other crops. Do not apply to rice under stress as phytotoxicity may result.

Do not add liquid nitrogen, zinc or surfactants to BOLERO7 8 EC.

\* \* \* \* \*

Follow all applicable directions, restrictions, and precautions on the EPA-registered label.

This labeling must be in the possession of the user at the time of pesticide application.

24(c) Registrant:

VALENT U.S.A. Corporation  
P.O. Box 8025  
Walnut Creek, CA 94596-8025

BOLERO7 - Reg. TM of Kumiai Chemical Industry Co. Ltd. for thiobencarb herbicide.