



FIFRA Section 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY IN THE STATES OF
CA, ID, IL, IN, KY, MI, MN, OH, OR, PA, VA, WA and WI

VENOM[®] I N S E C T I C I D E EPA Reg. No. 59639-135

FIFRA Section 2(ee) Recommendation
For Additional Pests in Grape

DIRECTIONS FOR USE

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. Always read and follow all label directions, restrictions and precautions when using any pesticide alone or in tank mix combinations. The most restrictive labeling applies when using a tank mix.

Venom[®] Insecticide is toxic to bees exposed to direct treatment or to residue on blooming crops and weeds. Do not apply *Venom* Insecticide or allow it to drift onto blooming plants if bees are actively foraging in the treated area.

This 2 (ee) bulletin contains directions for use which do not appear on the EPA-registered label of *Venom* Insecticide. Please refer to the production label for additional information.

GRAPE

CROP	PESTS	PRODUCT RATES	SPECIAL IINSTRUCTIONS
Grape	Glassy-Winged Sharpshooter Grape Berry Moth (first and second generation only) Leafhoppers Grape Mealybug Multi-colored Asian Lady Beetle Thrips	FOLIAR: 2 to 3 oz/A (0.088 to 0.132 lb ai/A)	Higher water volumes provide improved insect control. Begin applications when first pest activity is noticed or when insects reach threshold levels per State and County Extension Service recommendations. Repeat, as needed, to maintain control but not more often than every 14 days. For best results, time application before a damaging population becomes established.
	Glassy-Winged Sharpshooter Leafhoppers Mealybugs Grape Mealybug Vine Mealybug Obscure Mealybug <i>Phylloxera</i> spp. (suppression only) Thrips	SOIL: 5 to 6 oz/A (0.226 to 0.264 lb ai/A)	Under severe pest pressure, use the higher recommended rates. For Mealybug control, apply between budbreak and pea-berry size. The rate applied affects the length of control. Use the high rate where infestations occur later in crop development, or where pest pressure is continuous. <i>Venom</i> Insecticide may be tank mixed with other insecticides for better knock-down and/or improved control of pests.

Continued

NOTE: Regardless of application method, do not apply more than a total of 12 oz of *Venom* Insecticide (0.528 lb ai) per acre per season.

Foliar Application

- Apply with air or ground equipment in adequate water for uniform coverage (5 to 10 gal/A by air or 50 to 300 gal/A by ground).
- Do not apply *Venom* Insecticide within one (1) day of harvest.
- Do not apply more than a total of 6 oz of *Venom* Insecticide (0.264 lb ai) per acre per season by foliar.

Soil Application

- Make only one (1) soil application per season.
- Apply with ground equipment in adequate water for uniform coverage (10 to 100 gal/A).
- Do not apply *Venom* Insecticide within twenty-eight (28) days of harvest.
- Do not apply more than a total of 6.0 oz of *Venom* Insecticide (0.264 lb ai) per acre per season.
- Apply specified dosage in sufficient carrier volume to insure uniform application and incorporate into the soil using drip or trickle irrigation water. Apply towards the end of the irrigation run to ensure the product does not leach past the root zone.

Follow instructions carefully.

Follow all applicable directions, restrictions and precautions on the registered label.

This labeling must be in possession of the user at the time of pesticide application.

Please contact Valent U.S.A. Corporation at 800-6-VALENT (682-5368) to determine if this product is registered in your state.

THIS RECOMMENDATION EXPIRES ON DECEMBER 31, 2012

Registrant: Valent U.S.A. Corporation
P.O. Box 8025
Walnut Creek, CA 94596-8025

Venom is a registered trademark of Valent U.S.A. Corporation.