

FIFRA Section 2(ee) Recommendation



GROUP	3	INSECTICIDE
-------	---	-------------

RESTRICTED USE PESTICIDE
DUE TO TOXICITY TO FISH AND AQUATIC ORGANISMS.
For retail sale to and use only by Certified Applicators, or persons under their direct supervision,
and only for those uses covered by the Certified Applicator's certification.

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF NY



EPA Reg. No. 59639-35
 FIFRA Section 2(ee) Recommendation
 For use on Brown Marmorated Stink Bug

DIRECTIONS FOR USE

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. Always read and follow all label directions, restrictions and precautions when using any pesticide alone or in tank-mix combinations. The most restrictive labeling applies when using a tank mix.

RECOMMENDED FOR BROWN MARMORATED STINK BUG CONTROL

Pests	Crops	Danitol® 2.4 EC Spray Application Rates	Special Instructions
Brown Marmorated Stink Bug	Bushberries Cotton Cucurbits Head and Stem Brassica	10 2/3 to 16 fl oz/A (0.2 to 0.3 lb ai/A)	Apply <i>Danitol</i> 2.4 EC Spray as a foliar spray . Sufficient water volume should be used to provide complete and thorough coverage. Higher water volumes provide improved insect control. Insect control may be improved by the addition of a non-ionic surfactant to the spray solution according to manufacture's recommendation. Make applications beginning when first pest activity is noticed and repeat as needed. Under severe pest pressure, use the higher recommended rates. <i>Danitol</i> 2.4 EC Spray may be mixed with and/or alternated with commonly used insecticides to comply with local IPM programs. Continued on page 2
	Fruiting Vegetables Pea (Succulent)	10 2/3 fl oz/A (0.2 lb ai/A)	
	Grape Pome Fruit Stone Fruit Strawberry	10 2/3 to 21 1/3 fl oz/A (0.2 to 0.4 lb ai/A)	

Continued:

- Do not exceed 2 2/3 pt of *Danitol* 2.4 EC Spray (42 2/3 fl oz, 0.8 lb ai) per acre per season for all crops except bushberries and strawberry.
- Do not exceed more than two applications totaling 2 2/3 pt of *Danitol* 2.4 EC Spray (42 2/3 fl oz, 0.8 lb ai) per acre to the same planting in 12 consecutive months in strawberry.
- Do not exceed 2 pt of *Danitol* 2.4 EC Spray (32 fl oz, 0.6 lb ai) per acre per season in bushberries.
- Do not apply within twenty one (21) days of harvest for currant, cotton and grape.
- Do not apply within fourteen (14) days of harvest for pome fruit.
- Do not apply within seven (7) days of harvest for cucurbits, head and stem brassica and pea (succulent).
- Do not apply within three (3) days of harvest for bushberries, fruiting vegetables and stone fruit.
- Do not apply within two (2) days of harvest for strawberry.

APPLICATION INFORMATION AND PRECAUTIONS

- Always read and follow use directions, restrictions and precautions for all products. When tank mixing, the most restrictive labeling applies.
- Follow all applicable directions, restrictions and precautions on the EPA-registered labels.
- This labeling must be in the possession of the user at the time of pesticide application.
- It is a violation of Federal law to use these products in a manner inconsistent with their labeling.

Please contact Valent U.S.A. Corporation at 1-800-6-VALENT (682-5368) to determine if this product is registered in your state.

THIS RECOMMENDATION EXPIRES ON DECEMBER 31, 2012

Products That Work, From People Who Care[®] | www.valent.com | 800-6-VALENT (682-5368)

Registrant: Valent U.S.A. Corporation
P.O. Box 8025
Walnut Creek, CA 94596-8025

Danitol is a registered trademark of Sumitomo Chemical Company, Ltd.
Products That Work, From People Who Care is a registered trademark of Valent U.S.A. Corporation.