

FIFRA Section 2(ee) Recommendation



GROUP 3 FUNGICIDE

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF CA



EPA Reg. No. 59639-147

FIFRA Section 2(ee) Recommendation for Additional Pests in
Almond, Apricot, Nectarine and Peach

DIRECTIONS FOR USE

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA.

RECOMMENDED SPRAY DIRECTIONS FOR ALMOND					
Diseases	Application Rates		When to Apply	Special Use Instructions	Use Restrictions
	oz/A	GPA			
Alternaria Leaf Spot (<i>Alternaria alternata</i>)	2.5 to 3.5 (0.08 to 0.11 lb ai/A)	100 to 400 Aerial: Minimum 20 GPA	If monitoring indicates the presence of Alternaria Leaf Spot, begin late spring treatments about mid-April. In orchards with a history of the disease, treat in mid-to late- April and 2 to 3 weeks later.	<i>Quash</i> Fungicide should be used as part of an Integrated Pest Management (IPM) program. Apply as a foliar spray in sufficient water to obtain thorough coverage of blossoms, foliage and/or fruit.	<ul style="list-style-type: none"> Do not apply within 25 days of harvest Do not make more than 2 applications after petal fall. Do not make more than 2 sequential applications before switching to a non-group 3 fungicide for resistance management. Do not make more than 4 applications per season. Do not apply more than 14 oz/A per season.
Powdery mildew (<i>Podosphaera leucotricha</i>)	2.5 to 3.5 (0.08 to 0.11 lb ai/A)		Following brown rot/blossom blight schedule (i.e., applications during bloom), make additional applications on a 10 to 14 day interval until terminal growth ceases.		
Rust (<i>Tranzschelia discolor</i>)	3.5 oz/A (0.11 lb. ai/A)		Begin applications prior to disease development and continue at a 7 to 14 day interval throughout the season.		
Scab (<i>Cladosporium carpophilum</i>)	2.5 to 3.5 (0.08 to 0.11 lb ai/A)				

RECOMMENDED SPRAY DIRECTIONS FOR APRICOT, NECTARINE AND PEACH

Diseases	Application Rates		When to Apply	Special Use Instructions	Use Restrictions
	oz/A	GPA			
Powdery Mildew (<i>Podosphaera</i> spp.)	3.5 to 4.0 (0.11 to 0.125 lb ai/A)	100 to 400 Aerial: Minimum 20 GPA	Following brown rot/blossom blight schedule (i.e., applications during bloom), make additional applications on a 10 to 14 day interval until terminal growth ceases.	<i>Quash</i> Fungicide should be used as part of an Integrated Pest Management (IPM) program. Apply as a foliar spray in sufficient water to obtain thorough coverage of blossoms, foliage and/or fruit. Under high disease pressure the higher rate and shorter spray intervals should be used.	<ul style="list-style-type: none"> Do not apply within 14 days of harvest. Do not make more than 2 applications after petal fall. Do not make more than 2 sequential applications before switching to a non-group 3 fungicide for resistance management. Do not make more than 3 applications per season. Do not apply more than 12 oz/A per season.
Rust (<i>Tranzschelia discolor</i>)	3.5 oz/A (0.11 lb. ai/A)		Begin applications prior to disease development and continue at a 7 to 14 day interval throughout the season.		

This 2(ee) bulletin contains directions for use, which do not appear on the package label of *Quash* Fungicide.

Follow instructions carefully.

Follow all applicable directions, restrictions and precautions on the registered label.

Always read and follow all label directions, restrictions and precautions when using any pesticide alone or in tank mix combinations. The most restrictive labeling applies when using a tank mix.

This labeling must be in possession of the user at the time of pesticide application.

THIS RECOMMENDATION EXPIRES DECEMBER 31, 2012

Registrant: Valent U.S.A. Corporation
P.O. Box 8025
Walnut Creek, CA 94596-8025

Quash is a registered trademark of Valent U.S.A. Corporation