

FIFRA Section 2(ee) Recommendation



FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF
AL, AR, DE, GA, KY, MD, MS, NC, NJ, SC, TN & VA



VALOR[®] SX HERBICIDE TANK MIXED WITH DICAMBA, DICAMBA AND
2,4-D OR DICAMBA AND GLYPHOSATE FOR BURNDOWN AND
RESIDUAL CONTROL OF GLYPHOSATE-RESISTANT MARESTAIL
AND OTHER WINTER ANNUAL BROADLEAF WEEDS
EPA Reg. No. 59639-99
FIFRA SECTION 2(ee) RECOMMENDATION

DIRECTIONS FOR USE

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA.

USE VALOR TANK MIXED WITH DICAMBA, DICAMBA AND 2,4-D OR DICAMBA AND GLYPHOSATE FOR BURNDOWN AND RESIDUAL CONTROL OF GLYPHOSATE-RESISTANT MARESTAIL AND OTHER WINTER ANNUAL BROADLEAF WEEDS

Product	Product Rates	Weeds Controlled	Special Instructions
<i>Valor</i>	2 to 3 oz/A	Glyphosate-resistant marestail	Add a non-ionic surfactant at 1 qt/100 gal of spray solution.
plus			
Dicamba	8 fl oz/A	Refer to the <i>Valor</i> label for additional weeds controlled and sizes.	Apply in a minimum of 10 gal of spray solution per acre. Under high weed populations, a minimum of 15 to 20 gal of spray solution per acre should be applied.
or			
2,4-D + Dicamba	1 to 2 pt/A + 8 fl oz/A	Refer to 2,4-D, dicamba and glyphosate labels for weeds controlled and weed sizes.	
or			
Dicamba + Glyphosate	8 fl oz/A + 2 pt/A		

Use Restrictions for *Valor* plus dicamba, dicamba and 2,4-D or dicamba and glyphosate tank mix:

1. Always read and follow all label directions, restrictions and precautions when using any pesticide alone or in tank-mix combinations. **The most restrictive labeling applies when using a tank mix.**
2. The *Valor* plus dicamba and 2,4-D at 1 pt/A tank mix may be applied not less than 15 days prior to planting soybeans. If the 2,4-D rate is increased to 2 pt/A, then this tank mix may be applied not less than 30 days prior to planting soybeans. A minimum accumulation of 1 inch rainfall or overhead irrigation must also occur before soybeans can be planted. This interval must be observed prior to planting soybeans or crop injury may occur.
3. The *Valor* plus dicamba or dicamba plus glyphosate tank mix may be applied not less than 14 days prior to planting soybeans. A minimum accumulation of 1 inch rainfall or overhead irrigation must also occur before soybeans can be planted. This interval must be observed prior to planting soybeans or crop injury may occur.

This 2(ee) bulletin contains directions for use, which do not appear on the package label of *Valor* SX Herbicide. Follow instructions carefully.

Follow all applicable directions, restrictions and precautions on the EPA-registered labels.

It is a violation of Federal law to use these products in a manner inconsistent with its labeling.

THIS RECOMMENDATION EXPIRES DECEMBER 31, 2012

Valor is a registered trademark of Valent U.S.A. Corporation