

FIFRA Section 2(ee) Recommendation



FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF AL, AR, DE, GA, IA, IL, IN, KS, KY, LA, MD, MI, MS, NC, NE, NJ, OH, OK, PA, SC, TN, VA, WI



VALOR® XLT SOYBEAN HERBICIDE APPLIED EARLY PREPLANT BURNDOWN OR PREEMERGENCE FOLLOWED BY A POSTEMERGENCE APPLICATION OF WARRANT™ HERBICIDE OR METOLACHLOR CONTAINING PRODUCTS LABELED FOR POSTEMERGENCE USE IN SOYBEANS
FIFRA SECTION 2(ee) RECOMMENDATION

DIRECTIONS FOR USE

This recommendation is made as permitted under Section 2(ee) and has not been submitted to or approved by the EPA.

VALOR XLT SOYBEAN HERBICIDE APPLIED EARLY PREPLANT OR PREEMERGENCE FOLLOWED BY WARRANT HERBICIDE OR METOLACHLOR CONTAINING PRODUCTS LABELED FOR POSTEMERGENCE USE IN SOYBEANS.

BROADLEAF WEED SPECIES				
COMMON NAME	SCIENTIFIC NAME	ORGANIC MATTER	SOIL TYPE	RATES ¹
Carpetweed	<i>Mollugo verticillata</i>	Up to 5%	All Soil Types	Valor XLT Soybean Herbicide at 3 to 5 oz/A Followed by a postemergence application of: Warrant Herbicide or metolachlor containing products at labeled rates
Chickweeds				
Common	<i>Stellaria media</i>			
Mouseear	<i>Cerastium vulgatum</i>			
Dandelion	<i>Taraxacum officinale</i>			
Eclipta	<i>Eclipta prostrata</i>			
Eveningprimrose, Cutleaf	<i>Oenothera laciniata</i>			
Field Pennycress	<i>Thlaspi arvense</i>			
Florida Pusley	<i>Richardia scabra</i>			
Henbit	<i>Lamium amplexicaule</i>			
Lambsquarters, Common	<i>Chenopodium album</i>			
Little Mallow	<i>Malva parviflora</i>			
Marestail/Horseweed	<i>Conyza Canadensis</i>			
Mayweed/False Chamomile	<i>Matricaria maritime</i>			
Nightshades				
Black	<i>Solanum nigrum</i>			
Eastern Black	<i>Solanum ptycanthum</i>			
Hairy	<i>Solanum sarrachoides</i>			

¹ Refer to label for rates based on soil texture and target weed(s)

BROADLEAF WEED SPECIES				
COMMON NAME	SCIENTIFIC NAME	ORGANIC MATTER	SOIL TYPE	RATES ¹
Pigweeds		Up to 5%	All Soil Types	Followed by a postemergence application of: Warrant Herbicide or metolachlor containing products at labeled rates
Redroot	<i>Amaranthus retroflexus</i>			
Smooth	<i>Amaranthus hybridus</i>			
Spiny Amaranth	<i>Amaranthus spinosus</i>			
Tumble	<i>Amaranthus albus</i>			
Prickly Lettuce	<i>Lactuca serriola</i>			
Prickly Sida (Teaweed)	<i>Sida spinosa</i>			
Puncturevine	<i>Tribulus terrestris</i>			
Purslane, Common	<i>Portulaca oleracea</i>			
Radish, Wild	<i>Raphanus raphanistrum</i>			
Redmaids	<i>Calandrinia ciliata</i> var. <i>menziesii</i>			
Shepherd's-purse	<i>Capsella bursa-pastoris</i>			
Smallflower Morningglory	<i>Jacquemontia tamnifolia</i>			
Sowthistle, Prickly	<i>Sonchus asper</i>			
Spotted Spurge	<i>Euphorbia maculate</i>			
Venice Mallow	<i>Hibiscus trionum</i>			

¹ Refer to label for rates based on soil texture and target weed(s).

1. Do not graze animals on green forage or stubble. Do not utilize hay or straw for animal feed or bedding.
2. Always read and follow all label directions when using any pesticide alone or in tank mix combinations. The most restrictive labeling applies when using a tank mix.

Follow all applicable directions, restrictions and precautions on the EPA-registered labels. This labeling must be in the possession of the user at the time of pesticide application.

It is a violation of Federal law to use these products in a manner inconsistent with its labeling.

THIS RECOMMENDATION EXPIRES DECEMBER 31, 2012

Valor is a registered trademark of Valent U.S.A. Corporation
Warrant is a trademark of Monsanto Technology, LLC.