



NipsIt INSIDE[®] Insecticide

TREATED RICE SEED

This seed has been treated with NipsIt INSIDE[®]
Insecticide containing 47.8% clothianidin.

CAUTION
DO NOT USE FOR FEED,
FOOD OR OIL PROCESSING
STORE AWAY FROM FEEDS AND
OTHER FOODSTUFFS
KEEP OUT OF REACH OF CHILDREN

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear long-sleeved shirt, long pants and waterproof gloves when handling treated seed.

FIRST AID

IF SWALLOWED:

Call a poison control center or doctor immediately for treatment advice.

Have a person sip a glass of water if able to swallow.

Do not induce vomiting unless told to do so by a poison control center or doctor.

Do not give anything to an unconscious person.

Have product container or label with you when calling a poison control center or doctor, or going for treatment.

PLANT BACK RESTRICTIONS

ROTATIONAL CROP PLANT-BACK INTERVALS

Crops Planted	Immediate Plant-back	30-Day Plant-back	8-Month Plant-back	12-Month Plant-back
Rice	Bulb Vegetables, Canola, Cereal Grains, Cotton, Fruiting Vegetables (including Cucurbits), Leafy Vegetables (including Brassica Vegetables), Rapeseed, Soybean, Root and Tuber Vegetables	Dry Beans, Grasses, Non-grass Animal Feeds	Sugarcane	Any crops without an earlier plant-back interval

NOTE:

- Dispose of all excess treated seed. Left over treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements. Do not contaminate water bodies when disposing of planting equipment washwaters.
- Dispose of seed packaging in accordance with local requirements.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading.
- Treated seed does not flow as freely as untreated seed. Make adjustments to compensate for this restricted seed flow to prevent lower than desired seeding rates.
- Do not plant more than 150 lb of *NipsIt INSIDE* treated rice seed per acre (a total of 0.11 lb ai Clothianidin/150 lb seed/A) per year.
- *NipsIt INSIDE* Insecticide is to be used as a rice seed treatment ONLY when rice seed is dry-seeded in drill planters.
- DO NOT use *NipsIt INSIDE* Insecticide as a rice seed treatment in areas where rice/crawfish aquaculture practices are in place, or in or near fish farm, shrimp, prawn, or crab pond or nursery operations.
- DO NOT follow with a spray or foliar treatment of another neonicotinoid insecticide. Follow approved guidelines established for insect resistance management by consulting your local extension pest management expert.
- Tail water from rice fields planted with *NipsIt INSIDE* Insecticide treated rice seed MAY NOT be used to irrigate crops.
- Only for use in AR (not to exceed 20K acres), LA (not to exceed 5K acres), MS (not to exceed 5K acres), TX (not to exceed 10K acres).

**USE RATE
Rice**

Insect Pests	<i>NipsIt INSIDE</i> Insecticide Use Rates fl oz/100 lbs seed	Further Use Instructions
Grape Colaspis Rice Water Weevil	1.92 (0.075 lb ai/100 lbs seed)	<i>NipsIt INSIDE</i> Insecticide provides early season protection against listed pests. Do not plant more than 150 lb of <i>NipsIt INSIDE</i> treated rice seed per acre (a total of 0.11 lb ai Clothianidin /150 lb seed /A) per year.



EPA Experimental Use Permit No. 59639-EUP-15

NipsIt INSIDE is a registered trademark of Valent U.S.A. Corporation

For **PRODUCT USE** information:
Call 1-800-6-VALENT (1-800-682-5368)
IN CASE OF EMERGENCY ONLY
Call 24 Hours A Day 1-800-892-0099

Valent U.S.A. Corporation
P.O. Box 8025
Walnut Creek, California 94596-8025

Form 1797-A