

**SUPPLEMENTAL LABEL**



FOR ORGANIC PRODUCTION

Active Ingredient:  
 Gibberellic Acid..... 4.0% w/w  
 Other Ingredients..... 96.0% w/w  
 Total..... 100.0% w/w

EPA Registration No.: 73049-15  
 EPA Est. No.: 33762-IA-oo1

**(For Use In AL, AR, AZ, DE, FL, GA, KS, KY, LA, MD, MO, MS, NC, NM, NV, OK, SC, TN, TX, VA and WV Only)**

**Expires: December 31, 2015**

This supplemental label expires on 12/31/15 and must be not used or distributed after this date.

**KEEP OUT OF REACH OF CHILDREN  
 WARNING**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. READ THE LABEL AFFIXED TO THE CONTAINER FOR ProGibb 4% PLANT GROWTH REGULATOR BEFORE APPLYING. USE OF ProGibb 4% PLANT GROWTH REGULATOR ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR ProGibb 4% PLANT GROWTH REGULATOR.**

**COTTON:**

<p><b>ProGibb 4%</b> plant growth regulator has been shown to help shorten the vegetative growth “lag” phase. This benefit reduces the time interval needed to develop optimum leaf area and plant height, thus maximizing the potential for earliness and improved yields.</p>			
USE	OBJECTIVE/ BENEFIT	RATE (fl oz/acre)	APPLICATION TIMING
On young cotton plants	Promote growth and increase seedling vigor	<p>1 to 6 fl oz (30 to 180 ml)</p> <p>Use higher rates (within the indicated range) when temperatures will likely average 75°F or less during the 14 days following application(s).</p>	<p>In-furrow application to seed, or as a foliar application from the cotyledon leaf stage through the 7 leaf/node stage. Repeat applications as needed to a maximum of 3 applications.</p> <p>Applying more often than necessary to achieve the desired height results in excessive vegetative growth.</p>
<p><b>Do not</b> apply <b>ProGibb 4%</b> to cotton plants that are under drought stress. If the cotton plants are under continuous stress, delay the application of <b>ProGibb 4%</b> until the stress is alleviated and the plants are beginning to recover.</p> <p>Avoid drift or accidental application to other crops.</p>			

### **Mixing Instructions**

Fill the treatment tank with half of the final tank mix volume. Add the required amount of **ProGibb 4%** and mix thoroughly while adding water to the desired final volume. Dispose of any unused spray material at the end of the day.

### **Application Equipment**

Apply **ProGibb 4%** by aerial or ground spray equipment. As an aerial spray, use a spray system capable of producing a uniform pattern of medium to fine spray droplets at 10 gallons per acre (GPA). Apply no less than 3 GPA of total spray volume. Use low pressure ground sprayers equipped with boom and flat fan nozzles using 10 to 15 GPA spray volume.

**PLEASE CONTACT VALENT U.S.A. CORPORATION AT 800-6-VALENT (682-5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE.**

Copyright © 2012 by Valent U.S.A. Corporation

**ProGibb** is a registered trademark of Valent BioSciences Corporation.



Distributed by:  
**Valent U.S.A. Corporation**  
P.O. Box 8025  
Walnut Creek, CA 94596-8025  
[www.valent.com](http://www.valent.com)

Made in U.S.A.  
Form: 2012-PG4%-0010  
00000000