

Supplemental Label



GROUP	3	FUNGICIDE
-------	---	-----------



EPA Reg. No. 59639-147
(For New York Only)

QUASH™ FUNGICIDE FOR USE ON BUSHBERRIES (SUBGROUP 13-07B) AND TUBEROUS AND CORM VEGETABLES (Subgroup 1C)

This supplemental label expires on December 31, 2016 and must not be used or distributed after this date.

This supplemental label indicates removal of the statement "Do not use with an adjuvant or surfactant" from listed crop use directions, and thus supersedes any container or previous supplemental labels where this statement appears.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS, DIRECTIONS AND RESTRICTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. READ THE LABEL AFFIXED TO THE CONTAINER FOR *QUASH* FUNGICIDE BEFORE APPLYING. USE OF *QUASH* FUNGICIDE ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR *QUASH* FUNGICIDE.

**CROP SPECIFIC DIRECTIONS, RESTRICTIONS AND LIMITATIONS
BUSHBERRIES (Subgroup 13-07B)**

Aronia Berry; Blueberry, Highbush; Blueberry, Lowbush; Buffalo Currant; Chilean Guava; Cranberry, highbush; Currant, Black; Currant, Red; Elderberry; European Barberry; Gooseberry; Honeysuckle, (edible); Huckleberry; Jostaberry; Juneberry (Saskatoon Berry); Lingonberry; Native Currant; Salal; Sea Buckthorn; Cultivars, Varieties and/or hybrids of these					
Disease	Application Rates		When to Apply	Special Use Instructions	Use Restrictions
	oz/A	GPA			
Alternaria Leaf Spot and Fruit Rot (<i>Alternaria tenuissima</i>)	2.5 (0.078 lb ai/A)	Ground: Minimum 20 GPA Aerial: Minimum 10 GPA	Apply when conditions favor disease development and prior to infection. Continue applications on a 7 to 14-day interval.	Use <i>Quash</i> Fungicide as part of an Integrated Pest Management (IPM) program. Use a non-group 3 fungicide, with activity on the target disease, in alternation with <i>Quash</i> Fungicide. Apply as a foliar spray in sufficient water to obtain through coverage of blossoms, foliage, and/or fruit.	<ul style="list-style-type: none"> • Do not apply within 7 days of harvest. • Do not make more than 3 applications per season. • Do not apply more than 7.5 oz/A per season. • Do not make more than 2 sequential applications.
Anthracoze Fruit Rot (Ripe Rot) (<i>Colletotrichum</i> spp.)					
Botryosphaeria Stem Canker and Blight (<i>Botryosphaeria</i> spp.)					
Botrytis Blight and Fruit Rot (<i>Botrytis cinerea</i>)					
Exobasidium Fruit and Leaf Spot (<i>Exobasidium vaccinii</i>)					
Leaf Rust (<i>Pucciniastrum vaccinii</i>)					
Mummy Berry (<i>Monilinia vaccinii corymbosi</i>)					
Phomopsis Canker, Leaf Spot, Twig Blight and Fruit Rot (<i>Phomopsis vaccinii</i>)					
Powdery Mildew (<i>Microsphaera vaccinii</i>)					
Septoria Leaf Spot and Stem Canker (<i>Septoria albopunctata</i>)					

**CROP SPECIFIC DIRECTIONS, RESTRICTIONS AND LIMITATIONS
TUBEROUS AND CORM VEGETABLES (Subgroup 1C)**

Arracacha; Arrowroot; Artichoke, Chinese; Artichoke, Jerusalem; Canna, (edible); Cassava, (Bitter and Sweet); Chayote (root); Chufa; Dasheen (taro); Ginger; Leren; Potato; Sweet Potato; Tanier; Turmeric; Yam Bean; Yam, True					
Diseases	Application Rates		When to Apply	Special Use Instructions	Use Restrictions
	oz/A	GPA			
Black Dot (<i>Colletotrichum coccodes</i>)	2.5 to 4.0 (0.078 to 0.125 lb ai/A)	Ground: Minimum 10 Aerial: Minimum 5 GPA	Apply when conditions favor disease development and prior to infection. If conditions favor disease development, make additional applications at 7 to 10-day intervals.	Use <i>Quash</i> Fungicide as part of an Integrated Pest management (IPM) program. Apply as a foliar spray in sufficient water to obtain thorough coverage of plant.	<ul style="list-style-type: none"> • Do not make more than 4 applications/season. • Do not make more than 2 sequential applications. • Do not apply within 1 day of harvest. • Do not apply more than 16 oz/A per season.
Brown Spot (<i>Alternaria alternata</i>)					
Early Blight (<i>Alternaria solani</i>)					
Gray Mold (<i>Botrytis cinerea</i>) (suppression)					
Powdery Mildew (<i>Erysiphe cichoracearu</i>)					
White Mold (<i>Sclerotinia sclerotiorum</i>)	4.0 (0.125 lb ai/A)		Make first application prior to infection, generally at row closure and/or first bloom. Make second application 14 days later if conditions favor white mold development.		

PLEASE CONTACT VALENT U.S.A. CORPORATION AT 800-6-VALENT (682-5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE.

©2013 Valent U.S.A. Corporation

Quash is a trademark of Valent U.S.A. Corporation

Manufactured for
Valent U.S.A. Corporation
P. O. Box 8025
Walnut Creek, CA 94596-8025

Made in U.S.A.

Form 2013-QUA-0011