



ACTIVE INGREDIENT:

Bacillus thuringiensis, subsp. *kurstaki*, strain ABTS-351, fermentation solids, spores, and insecticidal toxins 2.3%

OTHER INGREDIENTS 97.7%

TOTAL 100.0%

Potency: 1600 Cabbage Looper Units of Potency per mg or 0.726 Billion Cabbage Looper Units/LB.

The percent active ingredient does not indicate product performance and potency measurements are not federally standardized.

EPA Reg. No. 73049-14

EPA Est. No. 33762-IA-001

List No. 05545

INDEX:

- 1.0 First Aid
- 2.0 Precautionary Statements
 - 2.1 Hazard to Humans (and Domestic Animals)
 - 2.2 Personal Protective Equipment (PPE)
 - 2.3 User Safety Recommendations
 - 2.4 Environmental Hazards
- 3.0 Directions for Use
- 4.0 Agricultural Use Requirements
- 5.0 Storage and Disposal
- 6.0 General Information
- 7.0 Notice
- 8.0 Application
- 9.0 Notice to User

KEEP OUT OF REACH OF CHILDREN

CAUTION

1.0

FIRST AID	
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 800-892-0099 (24 hours) for emergency medical treatment and/or transport emergency information. For all other information, call 800-6-VALENT (682-5368).	

2.0

PRECAUTIONARY STATEMENTS

2.1

HAZARD TO HUMANS (AND DOMESTIC ANIMALS)

CAUTION

Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse.

2.2

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Mixers/loaders and applicators must wear a dust/mist filtering respirator meeting NIOSH standards of at least N-95, R-95 or P-95. Repeated exposure to high concentrations of microbial proteins can cause allergic sensitization.

2.3

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

2.4

Environmental Hazards

Do not apply directly to water, or to where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters and rinsate.

This product must not be applied aerially within 1/4 mile of any habitats of endangered species or threatened lepidoptera. No manual application can be made within 300 feet of any threatened or endangered lepidoptera.

3.0

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

CONTINUED

4.0 AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Shoes plus socks
- Waterproof gloves

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. If recycling is not available dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

6.0 GENERAL INFORMATION

DiPel® 10G is a granular formulation of *Bacillus thuringiensis* variety *kurstaki*, designed specifically as a selective insecticide for control of certain leaf-eating caterpillars (Lepidoptera). For practical determinations of active ingredient, *DiPel* 10G can be expressed as a 10% formula of *DiPel* WP. The active ingredient in *DiPel* 10G is a stomach poison and therefore must be eaten by larvae to be effective. Climatic extremes such as heavy rain or strong wind shortly after treatment may act to remove *DiPel* 10G from the feeding zone making re-treatment necessary.

7.0 NOTICE

Local conditions may affect the use of *DiPel* 10G. Consult State Agricultural Extension or Experiment Station Specialists for specific recommendations related to local crop protection problems. Valent makes no warranty, express or implied, including the warranties of commercially acceptable quality and/or fitness for any particular purpose concerning this material, other than those which are contained on this label.

5.0 STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a cool, dry place.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

8.0 APPLICATION

Crop	Pests	Rate	Comments
Seed Corn, Field Corn, Popcorn and Sweet Corn	European Corn Borer, (<i>Ostrinia nubilalis</i>), First and Second Generation	10 lbs/acre	Ground or aerial equipment First Generation: Apply <i>DiPel</i> 10G directly to corn whorls. Treat seed corn when 25 percent of plants show shot hole feeding in the whorls and field corn when 35 percent of whorls show feeding. Larvae should be small and present at this time. Satisfactory control cannot be expected if treatment is delayed to higher injury levels or after larvae have begun stalk boring. Second Generation: Apply <i>DiPel</i> 10G when a field count shows 50 egg masses per 100 plants and the first hatch is occurring or follow your State Extension Service recommendation. Under conditions of intensive population pressure, a second application may be necessary. Ground or aerial equipment: Apply <i>DiPel</i> 10G directly to corn whorls. Treat sweet corn when 10-12 percent of plants show feeding in the whorls. Field corn should be treated when 75 percent of the plant whorls show feeding damage. Treatment thresholds vary for different types of corn so consult State Agricultural Extension specialists for specific recommendations. Under high pest pressure, a second application may be necessary.
	Southwestern Corn Borer (<i>Diatraea grandiosella</i>)	5-10 lbs/acre banded over-row with ground application	
	Fall Armyworm (<i>Spodoptera frugiperda</i>)	10 lbs/acre	
Tobacco	Tobacco Budworm (<i>Heliothis virescens</i>)	5-10 lbs/acre	Ground equipment: Apply <i>DiPel</i> 10G by hand or properly calibrated mechanical ground equipment.

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all of these factors when making decisions.

9.0 NOTICE TO USER

Seller makes no warranty, express or implied, of merchantability, fitness or otherwise concerning the use of this product other than as indicated on the label. User assumes all risks of use, storage or handling not in strict accordance with accompanying directions.

DiPel is a registered trademark of Valent BioSciences Corporation.

Products That Work, From People Who Care is a registered trademark of Valent U.S.A. Corporation.

© 2014 Valent BioSciences Corporation



Registered and Manufactured by:
Valent BioSciences Corporation
870 Technology Way
Libertyville, IL 60048

04-8054/R1 (Pkg. 04-7642/R2)

Distributed by Valent U.S.A. Corporation.
P.O. Box 8025
Walnut Creek, CA 94596-8025



For state registration and/or supplemental labels, please call or visit us online.

Products That Work, From People Who Care[®] | valent.com | 800-6-VALENT (682-5368)

Always read and follow label instructions.

© 2014 Valent BioSciences Corporation. All rights reserved. Printed in the U.S.A. 2014-D10G-0001 3/14