

**RESTRICTED USE PESTICIDE
DUE TO TOXICITY TO FISH AND AQUATIC ORGANISMS.
For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.**

Supplemental Label



**EPA Reg. No. 59639-35
(For Use in New York Only)**

DANITOL[®] 2.4 EC SPRAY FOR USE ON BARLEY

This supplemental label expires December 31, 2016 and must not be used or distributed after this date.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. READ THE LABEL AFFIXED TO THE CONTAINER FOR *DANITOL* 2.4 EC SPRAY BEFORE APPLYING. USE OF *DANITOL* 2.4 EC SPRAY ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR *DANITOL* 2.4 EC SPRAY

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

BARLEY		
PESTS	PRODUCT RATE FL OZ/A	SPECIAL INSTRUCTIONS
Aphids Cereal Leaf Beetle Thrips	10-2/3 (0.2 lb ai/A)	Apply as a ground application in a minimum of 5 gals of water per acre. Make a single application in the pre-boot stage. Resistance Management: It is generally recommended that no more than one <i>Danitol</i> 2.4 EC Spray application be made per season as part of a Resistance Management Program. Use non-pyrethroid products at other timings to control pests.
<ul style="list-style-type: none"> • Apply before the boot stage of growth(i.e., at or before Feekes Stage 9 (Zadoks Stage 4)). • Do not exceed 10-2/3 fl oz (0.2 lb ai) of <i>Danitol</i> 2.4 EC Spray per acre per season. 		

PLEASE CONTACT VALENT U.S.A. CORPORATION AT 800-6-VALENT (682-5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE.

©2015 Valent U.S.A. Corporation

Danitol is registered trademark of Sumitomo Chemical Company, Ltd.

Manufactured for
Valent U.S.A. Corporation
P. O. Box 8025
Walnut Creek, CA 94596-8025

Made in U.S.A.

2015-DAN-0012