

GROUP

3A

INSECTICIDE



Crop Protection EC 5.0_{II}



FOR ORGANIC PRODUCTION

- Provides rapid knockdown and kill of listed plant pests
- For use on growing crops
- Can be used on day of harvest

ACTIVE INGREDIENT:

Pyrethrins 5.00%

OTHER INGREDIENTS: 95.00%
100.00%

EPA Reg. No. 1021-1772
EPA Est. No. 1021-MN-2

**KEEP OUT OF REACH OF CHILDREN
CAUTION PRECAUCIÓN**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If swallowed:	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
If in eyes:	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on skin or clothing:	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

(continued)

FIRST AID (continued)

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-888-740-8712 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if absorbed through skin. Harmful if swallowed. Causes moderate eye irritation. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Avoid contact with skin, eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wear long-sleeved shirt and long pants, socks, shoes, and chemical-resistant gloves.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are made of any waterproof material. If you want more options, follow the instructions for category E on an EPA chemical-resistance category selection chart.

Mixers, loaders, applicators, and other handlers must wear the following:

- long-sleeved shirt,
- long pants,
- shoes and socks, and
- chemical-resistant gloves.

In addition to the above PPE, applicators using hand-held foggers in an enclosed area must wear a half-face, full-face, or hood-style NIOSH-approved respirator with:

- a dust/mist filtering cartridge (MSHA/NIOSH approval number prefix TC-21C), or
- a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G), or
- a cartridge or canister with any R, P or HE filter.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

When handlers use aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides 40 CFR 170.240 (d) (4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS:

Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic organisms, including fish and invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. This product may contaminate water through runoff. This product has a potential for runoff for several weeks after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR, Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas (that is permitted under Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water), is:

- Long-sleeved shirt and long pants;
- Chemical-resistant gloves, such as Barrier Laminate, Nitrile Rubber, Neoprene Rubber or Viton;
- Shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR, Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter until vapors, mists, and aerosols have dispersed, and the treated area has been thoroughly ventilated.

Do not enter or allow others to enter until sprays have dried.

INDOOR AREAS for use in and around:

Greenhouses

OUTDOOR AREAS for use in and around:

Nurseries

Outdoor Growing Crops

TO KILL THE FOLLOWING INSECTS:

Angoumois Grain Moths	Eriophyid Mites	Onion Maggots
Ants (excluding fire and Pharaoh ants)	European Pine Tip Moths	Orange Tortrix
Aphids	Fireworms	Pacific Flatheaded Borers
Apple Maggots	Flea Beetles	Pear Psyllids
Armored Scales	Fleas	Pepper Weevils
Armyworms	Forest Tent Caterpillars	Pink Bollworms
Artichoke Plume Moths	Fruit Flies	Potato Leafhoppers
Asian Citrus Psyllids	Fruittree Leafrollers	Potato Tuberworms
Asparagus Beetles	Fruitworms	Poultry Lice
Bagworms	Fungus Gnats	Proba Bugs
Bean Beetles	Garden Symphylan	Psyllids
Beet Armyworms	Glassy-winged Sharpshooters	Red Flour Beetles
Beetles	Gnats	Rice Weevils
Bermudagrass Mirids	Grain Mites	Saltmarsh Caterpillars
Billbugs	Grapeleaf Skeletonizers	Sawtoothed Grain Beetles
Blister Beetles	Grape Leafhoppers	Scales
Blow Flies	Grape Mealybugs	Sharpshooters
Boll Weevils	Grasshoppers	Sheep "Tick" or Ked
Boxelder Bugs	Greenbugs	Shore Flies
Branch and Twig Borers	Green Fruitworms	Shothole Borers
Cabbage Loopers	Green Peach Aphids	Silverfish
Cabbage Maggots	Greenhouse Thrips	Skippers
Cadellles	Gypsy Moths (adults & larvae)	Small Flying Moths
Cankerworms	Harlequin Bugs	Soft Scales
Carrot Weevils	<i>Heliothis</i> spp.	Southern Chinch Bugs
Caterpillars	Horn Flies	Sowbugs
Chalcids	Hornworms	Spider Beetles
Cheese Skippers	House Flies	Springtails
Cherry Fruit Flies	Imported Cabbageworms	Squash Bugs
Cigarette Beetles	Indian Meal Moths	Stable Flies
Clover Mites	Japanese Beetles	Stink Bugs
Clover Weevils	Katydid	Tarnished Plant Bugs
Cockroaches	Lace Bugs	Thrips
Codling Moths	Leaffooted Plant Bugs	Ticks
Colorado Potato Beetles	Leafhoppers	Tobacco Moths
Confused Flour Beetles	Leafminers	Tomato Budworms
Corn Earworms	Leafrollers	Tomato Bugs
Crane Flies	Leaf tiers	Tomato Fruitworms
Crickets	Loopers	Tomato Hornworms
Cross-striped Cabbageworms	Lygus	Tomato Pinworms
Cucumber Beetles	Maggots	Tomato Russet Mites
12-spotted Cucumber Beetles	Mealybugs	Tussock Moths
Dark Mealworms	Mediterranean Flour Moths	Vine Mealybugs
Darkling Beetles (lesser mealworms)	Mexican Bean Beetles	Vinegar Flies
Diamondback Larvae and Moths	Midges	Webworms
Driedfruit Beetles	Millipedes	Weevils
Drugstore Beetles	Mites	Western Yellowstriped Armyworms
Earwigs	Mole Crickets	Whiteflies
Eastern Tent Caterpillars	Moths	Yellow Mealworms
Elm Leaf Beetles	Mushroom Flies	
	Navel Orangeworms	
	Olive Fruit Flies	

GENERAL PRECAUTIONS AND RESTRICTIONS

Do not apply this product in a way that will contact workers or other persons, either directly or through drift.

Only protected handlers may be in the area during application.

Not for use in outdoor metered release devices.

Keep unprotected persons out of treated areas until sprays have dried.

Allow spray to dry before allowing adults, children or pets on treated areas.

Do not wet plants to the point of runoff or drip.

Do not apply more than 10 times per season.

Do not reapply within 3 days except under extreme pest pressure.

In case of extreme pest pressure, do not reapply within 24 hours.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Not for use in outdoor residential misting systems.

Do not apply when food, feed, and/or water is present.

Not for indoor use except in greenhouses.

Spray Drift Management for Agricultural Crops

Avoiding spray drift at the application site is the responsibility of the applicator and the grower. The interactions of many equipment and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

Do not apply at wind speeds greater than 10 mph at the application site.

Do not make any type of application into temperature inversions.

Apply as a medium or coarser spray (ASABE standard 572).

Additional Requirements for Aerial Applications:

Do not release spray at a height greater than 10 feet above the ground or crop canopy.

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

Aerial applicators must consider flight speed and nozzle orientation in determining droplet size.

When applications are made with a cross-wind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

Additional requirements for ground applications:

Do not release spray at a height greater than 4 feet above the ground or crop canopy.

Additional requirements for airblast applications:

Direct sprays into the canopy.

Turn off outward pointing nozzles at row ends and when spraying outer rows.

PESTICIDE RESISTANCE MANAGEMENT

PyGanic® Crop Protection EC 5.0_{II} contains a Group 3A insecticide. Insect biotypes with acquired resistance to Group 3A may eventually dominate the insect population if Group 3A insecticides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by PyGanic® Crop Protection EC 5.0_{II} or other Group 3A insecticides.

To delay insecticide resistance consider:

- Avoiding the consecutive use of PyGanic® Crop Protection EC 5.0_{II} or other Group 3A insecticides that have a similar target site of action, on the same insect species.
- Using tank mixtures or premixes with insecticides from a different target site of action Group as long as the involved products are all registered for the same use and have different sites of action.
- Basing insecticide use on a comprehensive IPM program.
- Monitoring treated insect population for loss of field efficacy.
- Contacting your local extension specialist, certified crop advisors, and/or manufacturer for insecticide resistance management and/or IPM recommendations for the specific site and resistance pest problems.

GENERAL USE DIRECTIONS

IMPORTANT: It is recommended that the final spray mix be buffered to a pH of 5.5-7.0. Outside of this range pyrethrins can degrade and the product will lose effectiveness.

To avoid possible harm to honey bees and other beneficial insects, it is advisable to apply when the non-target insects are least active.

Pyrethrins degrade rapidly in sunlight. To ensure the best performance, coverage is key. Use spray equipment that will provide the best coverage and direct contact with as many insects as possible.

Kills only at time of application. Listed insect pests must be directly contacted to be effective.

TANK MIX: This product may be tank mixed with other insecticides, acaricides, fungicides, adjuvants, and wetting agents. This application should conform to accepted use precautions and directions for both products. Tank mix applications must be made in accordance with the more restrictive of label limitations and precautions. No label application rates may be exceeded. This product cannot be mixed with any product with label prohibitions against such mixing.

Prior to tank mixing, a compatibility test should be

conducted using the proper proportions of products and water to ensure the physical compatibility of the mixture.

IMPORTANT NOTE: Plant safety is an important consideration when using insecticides in a greenhouse. However, it is not possible to evaluate the phytotoxicity of PyGanic® Crop Protection EC 5.0_{II} on numerous plant varieties that may react differently to insecticides in different growth stages or under varying environmental conditions. Before making widespread applications of PyGanic® Crop Protection EC 5.0_{II} or tank mix combinations, treat a limited number of plants and observe for phytotoxicity over a 10-day period.

DIRECTIONS FOR APPLICATION THROUGH IRRIGATION SYSTEMS (CHEMIGATION):

Apply this product only through sprinkler (including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move) irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for the operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the pump motor stops. The irrigation line or water pump must include a functional pressure valve, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

System must be a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Constant agitation must be maintained in the chemical supply tank during the entire period of insecticide application. Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of more dilute suspension per unit of time.

GROWING CROPS (OUTDOORS AND IN GREENHOUSES AND NURSERIES):

Apply 4.5 to 17 fl. oz. per acre by ground equipment in sufficient water for thorough coverage. This product may be applied by air at the rate of 4.5 to 17 fl. oz. per acre in a minimum of 2 gallons of water per acre for field crops and in a minimum of 10 gallons of water per acre for orchard crops. Mix only enough for immediate use. Spraying should begin when insects first appear. Do not wait until plants are heavily infested. Repeat as needed but not more than once daily.

PyGanic® Crop Protection EC 5.0_{II} may be used on most crops because its active ingredient is exempt from tolerances when applied to growing crops. The crop-grouping scheme used on this label was devised by the Environmental Protection Agency to expedite minor use pesticide registration. Each crop grouping on this label contains the phrase "including, but not limited to," and then lists a number of crops in each group. This wording allows the use of PyGanic® Crop Protection EC 5.0_{II} on crops that may not be specifically listed on this label (providing that the group to which the crop belongs is listed).

Maximum application rates:

Field and Orchard – 0.05 lbs. ai/acre or 17 fl. oz. of PyGanic® Crop Protection EC 5.0_{II} per acre (0.0012 lb. ai/1,000 sq. ft.)

Greenhouse space sprays – 0.0014 lbs. ai/acre.

ROOT AND TUBER VEGETABLES: Including: Arracacha, Arrowroot, Purple Arrowroot, Japanese Artichoke, Jerusalem Artichoke, Garden Beets, Sugar Beets, Edible Burdock, Edible Canna, Carrots, Cassava (bitter or sweet), Celeriac (Celery root), Chayote (root), Chervil (turnip rooted), Chicory, Chufa, Dasheen, Ginger, Ginseng, Horseradish, Leren, Parsley (turnip rooted), Parsnip, Potato, Radish, Japanese Radish (Daikon), Rutabaga, Salsify, Black Salsify, Spanish Salsify, Sweet Potato, Tanier, Turmeric, Turnip, Yam (true), Yam Bean.

LEAVES OF ROOT AND TUBER VEGETABLES: Including: Garden Beet, Sugar Beet, Edible Burdock, Carrot, Cassava (bitter or sweet), Celeriac (Celery root), Chervil (Turnip rooted), Chicory, Dasheen (Taro), Parsnip, Radish, Japanese Radish (Daikon), Rutabaga, Black Salsify, Sweet Potato, Tanier, Turnip, Yam (true).

BULB VEGETABLES: Including: Garlic, Great-headed Garlic, Leek, Onion (bulb and green) Welch, Shallot.

LEAFY VEGETABLES: Including: Amaranth (Leafy Amaranth, Chinese Spinach, Tampala), Arugula, Cardoon, Celery, Chinese Celery, Celtuce, Chervil, Corn Salad, Chrysanthemum (edible-leaved), Chrysanthemum (garland), Cress (garden, water), Upland Cress (Yellow Rocket, Winter Cress), Dandelion, Dock (Sorrel), Endive (Escarole), Fennel (Florence), Lettuce (head and leafy), Orach, Parsley, Purslane (garden & winter), Radicchio, Rhubarb, Spinach, Fine Spinach (Malabar, Ceylon), Spinach (New Zealand), Swiss Chard.

BRASSICA (COLE) LEAFY VEGETABLES: Including: Broccoli, Chinese Broccoli (Gai Lan), Broccoli Raab (Rapini), Brussels Sprouts, Cabbage, Chinese Cabbage (Bok Choy), Chinese Cabbage (Napa), Chinese Mustard Cabbage (Gai Choy), Cauliflower, Cavallo Broccoli, Collards, Kale, Kohlrabi, Mizuna, Mustard Greens, Mustard Spinach, Rape Greens.

LEGUME VEGETABLES (SUCCULENT OR DRIED): Including: Adzuki Beans, Field Beans, Kidney Beans, Lima Beans, Moth Beans, Mung Beans, Navy Beans, Pinto Beans, Rice Beans, Runner Beans, Snap Beans, Tepary Beans, Urd Beans, Wax Beans, Asparagus Beans, Blackeyed Peas, Catjang, Chinese Long Beans, Cowpeas, Crowder Peas, Southern Peas, Yard-long Beans, Broad Beans, (Fava Beans), Chick Peas (Garbanzo Beans), Guar, Jack Beans (Sword Bean), Lablab Beans (Hyacinth Bean), Lentils, Peas (Garden Peas, Field Peas, Sugar Snap Peas, English Peas, Snow Peas), Pigeon Peas, Soybeans, Sweet Lupin Beans, White Lupin Beans, White Sweet Lupin, Sword Beans.

FOLIAGE OF LEGUME VEGETABLES: Including: Plant part of any legume vegetable included in the legume vegetable group that will be used as animal feed including any variety of Beans, Field Peas, Soybeans.

FRUITING VEGETABLES: Including: Eggplant, Ground Cherry, Okra, Pepinos, Peppers (Bell Peppers, Chili Peppers, Cooking Peppers, Pimentos, Sweet Peppers), Tomatillos, Tomatoes.

CUCURBIT VEGETABLES: Including: Balsam Apple, Balsam Pear (Bitter Melon), Chayote (fruit), Chinese Wax Gourd (Chinese Preserving Melon), Chinese Cucumber, Citron Melon, Cucumber, Gherkin, Edible Gourds, Melons (including hybrids, Cantaloupe, Casaba, Golden Pershaw Melon, Crenshaw, Honeydew Melons, Honey Balls, Mango Melon, Muskmelon, Persian Melon, Pineapple Melon, Santa Claus Melon, Snake Melon), Pumpkin, Squash (summer & winter), Watermelon (including hybrids).

CITRUS FRUITS: Including: Calamondin, Citrus Citron, Citrus Hybrids, Grapefruit, Kumquats, Lemons, Limes, Mandarin (Tangerine), Orange (sweet & sour), Pummelo, Satsuma Mandarin, (*Citrus* spp. includes Chironja, Tangelos, Tangors).

POME FRUITS: Including: Apple, Crab Apple, Loquat, Mayhaw, Pear, Oriental Pear, Quince.

STONE FRUITS: Including: Apricot, Cherry (sweet & sour), Nectarine, Peach, Plum, Prune, Chickasaw Plum, Damson Plum, Japanese Plum, Plumcot.

SMALL FRUITS AND BERRIES: Including: Blackberry, Blueberry, Cranberry, Currant, Dewberry, Elderberry, Gooseberry, Grape, Huckleberry, Loganberry, Olallieberry, Raspberry (black & red), Strawberry, Youngberry.

TREE NUTS: Including: Almond, Beech Nut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (Hazelnut), Hickory Nut, Macadamia Nut (Bush Nut), Pecan, Pistachio, Walnut, Black and English (Persian).

ORIENTAL VEGETABLES: Including: Acerola, Atemoya, Balsam Pear (bitter melon), Carambola, Japanese Artichoke, Chinese Broccoli (Gai Lan), Chinese Cabbage (Bok Choy, Napa), Chinese Mustard Cabbage (Gai Choy), Dasheen, Ginger, Ginseng, Chinese Longbeans, Mung Beans, Citron Melon, Japanese Radish (Daikon), Chinese Spinach, Chinese Waxgourd, Cilantro, Citron Melon, Rambutan, Water Chestnut.

SUBTROPICAL FRUITS: Including: Avocado, Banana, Carob, Barbados Cherry, Cherimoya, Dates, Durian (Jackfruit), Feijoa, Figs, Guava, Kiwifruit, Lychee, Mango, Papaya, Passion Fruit, Persimmon, Pineapple, Pomegranate.

CEREAL GRAINS: Including: Barley, Buckwheat, Corn (sweet and field), Millet, Proso, Oats, Pearl Millet, Popcorn, Rice, Rye, Sorghum (Milo), Teosinte, Triticale, Wheat, Wild Rice.

FORAGE, FODDER AND STRAW OF CEREAL GRAINS: Including: Barley, Buckwheat, Corn (sweet and field), Millet, Proso, Oats, Pearl, Popcorn, Rice, Rye, Sorghum (Milo), Teosinte, Triticale, Wheat, Wild Rice.

GRASSES FOR SEED, FORAGE, FODDER AND HAY: Including: Any grass (Gramineal family, green or cured, except sugarcane and those listed in the cereal grains group) Pasture and Range Grasses, Grasses grown for hay and silage, Bermudagrass, Bluegrass, Bromegrass, Fescue, Oat Hay, Orchard Grass, Forage Sorghum, Sudan Grass, Timothy.

NON-GRASS ANIMAL FEEDS: Including: Alfalfa, Velvet Beans, Clover, Kudzu, Lespedeza, Lupine, Sainfoin, Trefoil, Crown Vetch, Milk Vetch.

HERBS AND SPICES: Including: Allspice, Angelica, Anise (Anise seed), Star Anise, Annatto (seed), Balm (Lemon balm), Basil, Borage, Burnet, Capers Buds, Caraway, Black Caraway, Cardamom, Cassia Bark, Cassia Buds, Catnip, Celery Seed, Chamomile, Chervil (dried), Chicory, Chive, Chinese Chive, Cinnamon, Clary, Clove Buds, Coriander (Cilantro or Chinese Parsley leaf), Coriander (Cilantro seed), Costmary, Culantro (leaf), Culantro (seed), Cumin, Curry (leaf), Dill (Dill Weed), Dill (seed), Fennel (common), Florence Fennel (seed), Fenugreek, Grains of Paradise, Horehound, Hyssop, Juniper Berry, Lavender, Lemongrass, Lovage (leaf), Lovage (seed), Mace, Mari-

gold, Marjoram (includes sweet or annual Marjoram, Wild Marjoram or Oregano, and Pot Marjoram), Mustard (seed), Nasturtium, Nutmeg, Oregano, Mint, Paprika, Parsley (dried), Pennyroyal, Black Pepper, White Pepper, Poppy (seed), Rosemary, Rue, Saffron, Sage, Summer and Winter Savory, Sweet Bay (Bay leaf), Tansy, Tarragon, Thyme, Vanilla, Wintergreen, Woodruff, Wormwood.

ADDITIONAL CROPS: Including: Artichoke, Asparagus, Avocado, Coffee, Cotton, Hops, Jojoba, Mushroom, Okra, Olives, Peanuts, Pineapple, Rice, Safflowers, Sesame, Sugar Cane, Sunflower, Tea.

TO KILL ANTS (EXCLUDING FIRE AND PHAROAH ANTS) ON ORCHARD FLOORS AND VINEYARD FLOORS: Apply as a broadcast spray to the entire orchard or vineyard floor using ground spray equipment at 5 to 17 fl. oz. per acre in 25 or more gallons of water. For best results use the high rate for heavy infestations and the lower rate for light infestations.

Do not apply where weed growth or other obstructions would impede uniform coverage of the orchard or vineyard floor. Mow or chemically control weeds before the application. Foliar applications of PyGanic® Crop Protection EC 5.0_{II} may be made in addition to the orchard floor treatment.

USE ON GREENHOUSE, ORNAMENTAL FLOWER AND FOLIAGE PLANTS: Combine 16 to 32 fl. oz. with 100 gallons of water for applications with conventional sprayers or 1/4 - 1/2 of a fl. oz. per gallon of water for applications with handheld or backpack sprayers. Apply the diluted product at the rate of 3 gallons/1,000 sq ft.

Do not exceed maximum application rate of .0036 lbs. ai/1,000 sq ft or 1.18 fl. oz. of PyGanic® Crop Protection EC 5.0_{II}/1,000 sq. ft.

Do not apply more than 1 time per day.

FOR USE IN VEGETABLE AND TRUCK GARDENS: For commercial production of listed crops to kill listed insects dilute 1/4 to 1/2 fl. oz. of PyGanic® Crop Protection EC 5.0_{II} per gallon of water. Apply diluted solution with all types of agricultural sprayers, including compressed air sprayers or pump driven sprayers.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Keep this product in its tightly closed original container, when not in use. Store in a cool, dry (preferably locked) area that is inaccessible to children and animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Non-refillable container. Do not reuse or refill this container.

Triple rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Products That Work, From People Who Care is a registered trademark of Valent U.S.A. Corporation.

MGK and PyGanic are registered trademarks of McLaughlin Gormley King Company. All other marks are property of their respective holders.

Manufactured by:

MGK
8810 Tenth Avenue North
Minneapolis, MN 55427
Rev. 0116-0816

Information contained in this booklet is accurate at the time of printing. Since product testing is a continuous process, please read and follow the directions on the product label for the most current directions and precautionary statements.

Always check with your state to verify state registration status or call 800-6-VALENT (682-5368).



For state registration and/or supplemental labels, please call.

Products That Work, From People Who Care® | 800-6-VALENT (682-5368)

Always read and follow label instructions.

©2016 Valent U.S.A. Corporation. All rights reserved. Printed in the U.S.A.