

Supplemental Label



PRESIDIO[®] FUNGICIDE

EPA Reg. No. 59639-140
(Except California)

PRESIDIO[®] FUNGICIDE FOR USE ON BASIL, LISTED SUCCULENT BEANS AND HOPS

This supplemental label expires on March 23, 2021 and must not be used or distributed after this date.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. READ THE LABEL AFFIXED TO THE CONTAINER FOR *PRESIDIO* FUNGICIDE BEFORE APPLYING. USE OF *PRESIDIO* FUNGICIDE ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR *PRESIDIO* FUNGICIDE.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SPECIFIC REQUIREMENTS – All Crops

When to Apply: Begin application when crop and/or environmental conditions favor disease development. Make applications on a 7- to 14-day interval, depending on disease pressure, unless otherwise noted under crop specific use instructions. Under severe disease pressure or if rain is expected, use the higher labeled rate and shortest interval. Apply as a foliar spray in sufficient water to obtain thorough coverage. Chemigation is not recommended when the primary disease is downy mildew or other foliar diseases.

Resistance Management: *Presidio* Fungicide must be used as part of an Integrated Pest Management (IPM) program. When tank mixing is required in the crop specific use instructions, use with another labeled fungicide product with a different mode of action on the target pathogen in sufficient water to obtain thorough coverage.

Restrictions and Limitations

- **Maximum rate *Presidio* Fungicide per application:** 4 fl oz per acre
- **Maximum rate of *Presidio* Fungicide per year unless otherwise noted under crop specific use instructions:** 12 fl oz per acre
- Apply no more than 2 sequential applications of *Presidio* Fungicide before alternating with an effective fungicide from a different resistance management group unless otherwise noted under crop specific use instructions.
- Do not make more than 4 applications of *Presidio* Fungicide per acre per year unless otherwise noted under crop specific use instructions.
- **In Hawaii:** Apply *Presidio* Fungicide by foliar and chemigation (sprinkler irrigation) only.

Basil				
Diseases	Application Rates		Minimum Time from Last Application to Harvest (PHI)	Specific Use Instructions
	fl oz/A	GPA Spray Mixture		
Downy Mildew (<i>Peronosporabelbahrii</i>)	4 (0.125 lb ai/A)	Ground: Minimum 20 Aerial: Minimum 5	1 day	Make foliar applications on a 7- to 14-day schedule beginning when conditions are favorable, but prior to disease development.

Listed Succulent Beans asparagus bean; Chinese longbean; moth bean; runner bean; snap bean; wax bean; yardlong bean				
Diseases	Application Rates		Minimum Time from Last Application to Harvest (PHI)	Specific Use Instructions
	fl oz/A	GPA Spray Mixture		
Phytophthora Root Rot (<i>Phytophthora parasitica</i>)	4 (0.125 lb ai/A)	Ground: Minimum 30	0 day	Make foliar applications on a 7- to 14-day schedule beginning when conditions are favorable, but prior to disease development.
Phytophthora Blight (<i>Phytophthora capsici</i>)		Aerial: Minimum 5		

Hops				
Diseases	Application Rates		Minimum Time from Last Application to Harvest (PHI)	Specific Use Instructions
	fl oz/A	GPA Spray Mixture		
Downy Mildew (<i>Pseudoperonospora humuli</i>)	4 (0.125 lb ai/A)	Ground: Minimum 50 Aerial: Minimum 5	24 days	Make foliar applications on a 10- to 14-day schedule beginning when conditions are favorable, but prior to disease development.

PLEASE CONTACT VALENT U.S.A. LLC AT 800-6-VALENT (682-5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE.

©2018 Valent U.S.A. LLC

Presidio is a registered trademark of Valent U.S.A. LLC

Manufactured for:
Valent U.S.A. LLC
P.O. Box 8025
Walnut Creek, CA 94596-8025
Made in U.S.A.

2018-PRD-0012