

# Supplemental Label



## PRESIDIO<sup>®</sup> FUNGICIDE

EPA Reg. No. 59639-140  
(For Use in California Only)

### PRESIDIO<sup>®</sup> FUNGICIDE FOR USE ON BASIL AND HOPS

This supplemental label expires on March 23, 2021 and must not be used or distributed after this date.

**THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. READ THE LABEL AFFIXED TO THE CONTAINER FOR *PRESIDIO* FUNGICIDE BEFORE APPLYING. USE OF *PRESIDIO* FUNGICIDE ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR *PRESIDIO* FUNGICIDE.**

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### SPECIFIC REQUIREMENTS – All Crops

**When to Apply:** Begin application when crop and/or environmental conditions favor disease development. Make applications on a 7- to 14-day interval, depending on disease pressure, unless otherwise noted under crop specific use instructions. Under severe disease pressure or if rain is expected, use the higher labeled rate and shortest interval. Apply as a foliar spray in sufficient water to obtain thorough coverage. Chemigation is not recommended when the primary disease is downy mildew or other foliar diseases.

**Resistance Management:** *Presidio* Fungicide must be used as part of an Integrated Pest Management (IPM) program. When tank mixing is required in the crop specific use instructions, use with another labeled fungicide product with a different mode of action on the target pathogen in sufficient water to obtain thorough coverage.

#### Restrictions and Limitations

- **Maximum rate *Presidio* Fungicide per application:** 4 fl oz per acre
- **Maximum rate of *Presidio* Fungicide per year unless otherwise noted under crop specific use instructions:** 12 fl oz per acre
- Apply no more than 2 sequential applications of *Presidio* Fungicide before alternating with an effective fungicide from a different resistance management group unless otherwise noted under crop specific use instructions.
- Do not make more than 4 applications of *Presidio* Fungicide per acre per year unless otherwise noted under crop specific use instructions.

**SPECIFIC REQUIREMENTS – ALL CROPS**

<b>Basil</b>				
<b>Diseases</b>	<b>Application Rates</b>		<b>Minimum Time from Last Application to Harvest (PHI)</b>	<b>Specific Use Instructions</b>
	<b>fl oz/A</b>	<b>GPA Spray Mixture</b>		
Downy Mildew ( <i>Peronosporabelbahrii</i> )	4 (0.125 lb ai/A)	Ground: Minimum 20  Aerial: Minimum 5	1 day	Make foliar applications on a 7- to 14-day schedule beginning when conditions are favorable, but prior to disease development.

<b>Hops</b>				
<b>Diseases</b>	<b>Application Rates</b>		<b>Minimum Time from Last Application to Harvest (PHI)</b>	<b>Specific Use Instructions</b>
	<b>fl oz/A</b>	<b>GPA Spray Mixture</b>		
Downy Mildew ( <i>Pseudoperonospora humuli</i> )	4 (0.125 lb ai/A)	Ground: Minimum 50  Aerial: Minimum 5	24 days	Make foliar applications on a 10- to 14-day schedule beginning when conditions are favorable, but prior to disease development.

**PLEASE CONTACT VALENT U.S.A. LLC AT 800-6-VALENT (682-5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE.**

©2018 Valent U.S.A. LLC

*Presidio* is a registered trademark of Valent U.S.A. LLC

Manufactured for:  
Valent U.S.A. LLC  
P.O. Box 8025  
Walnut Creek, CA 94596-8025  
Made in U.S.A.

2018-PRD-0013