

Supplemental Label



PRESIDIO[®] FUNGICIDE

EPA Reg. No. 59639-140
For Use in Indiana, Maryland, West Virginia and Wisconsin only

PRESIDIO[®] FUNGICIDE FOR USE ON TOBACCO

This supplemental label expires on January 31, 2020 and must not be used or distributed after this date.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. READ THE LABEL AFFIXED TO THE CONTAINER FOR *PRESIDIO* FUNGICIDE BEFORE APPLYING. USE OF *PRESIDIO* FUNGICIDE ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR *PRESIDIO* FUNGICIDE.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SPECIFIC REQUIREMENTS				
Restrictions and Limitations				
<ul style="list-style-type: none"> • Maximum rate <i>Presidio</i> Fungicide per application: 4 fl oz per acre. • Maximum rate of <i>Presidio</i> Fungicide per year unless otherwise noted under crop specific use instructions: 12 fl oz per acre. 				
Diseases	Application Rates		Minimum Time from Last Application to Harvest (PHI)	Specific Use Instructions
	fl oz/A	GPA Spray Mixture		
Black Shank* (<i>Phytophthora nicotianae</i>)	4 (0.125 lb ai/A)	Ground: Minimum 20	N/A	<p>Resistance Management Do not make sequential applications of <i>Presidio</i> Fungicide; alternate with a fungicide which has a different mode of action (different FRAC group) against the target pathogen.</p> <p>Do not make more than 2 soil spray applications per year.</p> <p>Do not make more than 3 total applications of <i>Presidio</i> Fungicide per year.</p> <p>Minimum treatment interval: 14 days.</p> <p>Post-transplant program If a fungicide with a different mode of action is used at or immediately after transplanting, one application of <i>Presidio</i> Fungicide can be made at either first cultivation or layby (last cultivation). Direct nozzles to cover soil beneath lower leaves; incorporate immediately with cultivator.</p>

*For best control of black shank with *Presidio* Fungicide, use with tobacco varieties that have moderate-to-high resistance to the black shank pathogen. Consult your local Cooperative Extension Service office or university specialist for information on variety selection.

Diseases	Application Rates		Minimum Time from Last Application to Harvest (PHI)	Specific Use Instructions
	fl oz/A	GPA Spray Mixture		
Blue Mold (<i>Peronospora tabacina</i>)	4 (0.125 lb ai/A)	Ground minimum: 20 Aerial minimum: 10	7 days	<p>Resistance Management Foliar applications of <i>Presidio</i> Fungicide must be tank mixed with a labeled rate of another fungicide active against the target pathogen, but with a different mode of action.</p> <p>Do not make more than 2 foliar applications per year.</p> <p>Do not make more than 3 total applications of <i>Presidio</i> Fungicide per year.</p> <p>Apply as a foliar spray prior to disease onset or at first indication that blue mold is in the area.</p> <p>Minimum retreatment interval: 7 days</p>

PLEASE CONTACT VALENT U.S.A. LLC AT 800-6-VALENT (682-5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE.

©2018 Valent U.S.A. LLC

Presidio is a registered trademark of Valent U.S.A. LLC

Manufactured for:
Valent U.S.A. LLC
P.O. Box 8025
Walnut Creek, CA 94596-8025
Made in U.S.A.

2018-PRD-0015
SAL20180612