

Supplemental Label



CHATEAU®

HERBICIDE SW

EPA Reg. No. 59639-99

(For Use in Arizona, Colorado, Hawaii, Idaho, Nebraska, Oregon and Washington)

CHATEAU® HERBICIDE SW USE IN DRY BEAN FOR WEED SUPPRESSION

Dried cultivars of bean (*Lupinus*); bean (*Phaseolus*) (includes field bean, kidney bean, lima bean (dry), navy bean, pinto bean, tepary bean); bean (*Vigna*) (includes adzuki bean, blackeyed pea, catjang, cowpea, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean); broad bean (dry); chickpea; guar; lablab bean and lentil

This supplemental label expires on August 31, 2019 and must not be used or distributed after this date.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. READ THE LABEL AFFIXED TO THE CONTAINER FOR CHATEAU HERBICIDE SW BEFORE APPLYING. USE OF CHATEAU HERBICIDE SW ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR CHATEAU HERBICIDE SW.

RESTRICTIONS AND LIMITATIONS

- Do not apply more than 1.5 oz of *Chateau* Herbicide SW per acre per application.
- Do not make more than 1 application of *Chateau* Herbicide SW per acre per year.
- Do not apply more than 1.5 oz of *Chateau* Herbicide SW per acre per year.

Many weather related factors, including high wind, splashing or heavy rains or cool conditions at or near crop emergence, may result in dry bean injury in fields treated with *Chateau* Herbicide SW. On occasion this has resulted in a delay in maturity. Users assume these risks before using *Chateau* Herbicide SW.

TIMING TO DRY BEAN

Chateau Herbicide SW may be applied to dry beans within 2 days after planting for the preemergence suppression of the weeds listed in Table A, Weeds Suppressed by Residual Activity of *Chateau* Herbicide SW at 1.5 oz/A. Tank mix *Chateau* Herbicide SW with other labeled herbicides for broad spectrum weed control.

TIMING TO WEEDS

Chateau Herbicide SW may be applied to dry beans prior to planting or preemergence (after planting). Preemergence application of *Chateau* Herbicide SW must be made within 2 days after planting and prior to dry bean emergence. To avoid severe crop injury, do not apply to dry beans after beans begin to crack or have emerged.

Preplant incorporation (PPI) applications may result in reduced weed control.

ADDITIONAL RESIDUAL GRASS CONTROL

Chateau Herbicide SW can be tank mixed with pendimethalin for additional grass control.

Table A. Weeds Suppressed by Residual Activity of *Chateau* Herbicide SW at 1.5 oz/A

COMMON NAME	SCIENTIFIC NAME	ORGANIC MATTER	CHATEAU HERBICIDE SW RATE
Lambsquarters, Common	<i>Chenopodium album</i>	Up to 5%	1.5 oz/A
Mustard, Wild	<i>Brassica kaber</i>		
Nightshades			
Black	<i>Solanum nigrum</i>		
Eastern Black	<i>Solanum ptycanthum</i>		
Hairy	<i>Solanum sarrachoides</i>		
Palmer Amaranth	<i>Amaranthus palmeri</i>		
Pigweeds			
Redroot	<i>Amaranthus retroflexus</i>		
Smooth	<i>Amaranthus hybridus</i>		
Spiny Amaranth	<i>Amaranthus spinosus</i>		
Tumble	<i>Amaranthus albus</i>		
Prickly Lettuce (China Lettuce)	<i>Lactuca serriola</i>		
Radish, Wild	<i>Raphanus raphanistrum</i>		

PLEASE CONTACT VALENT U.S.A. LLC AT 800-6-VALENT (682-5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE.

©2018 Valent U.S.A. LLC

Chateau is a registered trademark of Valent U.S.A. LLC

Manufactured for:

Valent U.S.A. LLC

P.O. Box 8025

Walnut Creek, CA 94596-8025

Made in U.S.A.

2018-CHSW-0016

SAL20180309