

# FIFRA Section 2(ee) Recommendation



FOR DISTRIBUTION AND USE ONLY IN THE STATES OF  
MICHIGAN, MINNESOTA, NORTH DAKOTA and WISCONSIN

## **BELAY<sup>®</sup>** INSECTICIDE

EPA Reg. No. 59639-150

### **FOR IMPREGNATION OF BELAY<sup>®</sup> INSECTICIDE ON DRY BULK FERTILIZER FOR USE ON POTATO**

This FIFRA Section 2(ee) Recommendation Expires December 31, 2021.

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This 2(ee) recommendation contains directions for use, which do not appear on the package label and must be in the possession of the user at the time of insect growth regulator application. Follow all applicable directions, restrictions and precautions on the EPA registered label when using *Belay* Insecticide alone or in tank mix combinations. The most restrictive labeling applies when using a tank mix.

#### **When Used on Potatoes**

*Belay* Insecticide can be impregnated or coated on dry bulk fertilizers. Many commonly used dry bulk fertilizers can be impregnated or coated with *Belay* with the exception of powdered limestone or fertilizers containing ammonium nitrate, potassium nitrate or sodium nitrate.

#### **Impregnate or Coat Dry Bulk Fertilizer with *Belay* Insecticide**

1. Mix required amount of *Belay* Insecticide in carrier to be sprayed onto fertilizer.
2. Utilize fertilizers that have the physical capacity of absorbing the carrier and product during the direct application into the fertilizer.
3. Apply *Belay* Insecticide directly to the dry fertilizer while blending covering the fertilizer only.
4. Allow treated fertilizer to blend for at least 5 minutes or until uniformly blended.

#### **Calculate the Amount of *Belay* Insecticide to be Used**

1. Determine rate from the *Belay* Insecticide label according to pest and pest pressure in the region. Use the rate of 12 fl oz/A when heavy infestations of target pest are expected.
2. Determine the amount of fertilizer to be treated (typically ranges from 300 to 1000 pounds per acre).
3. Use the following formula: (fl oz of *Belay* Insecticide) X (2000 pounds of fertilizer per acre)

#### **Field Application**

1. Apply the impregnated/coated fertilizer to the field as soon as possible after treating.
2. Accurate calibration and application will maximize uniform distribution and maximize performance.
3. An inconsistent and/or uneven impregnation/coating of fertilizer will result in poor product performance. Application of fertilizer coated/impregnated with *Belay* Insecticide needs to be delivered with properly calibrated equipment and uniform distribution pattern in the furrow during planting operations.

Refer to *Belay* Insecticide label for additional insects controlled.

**Please contact Valent U.S.A. LLC at 800-6-VALENT (682-5368) to determine if this product is registered in your state.**

*Belay* is a registered trademark of Sumitomo Chemical Company, Ltd.

Registrant: Valent U.S.A. LLC  
P.O. Box 8025  
Walnut Creek, CA 94596-8025