



TREATED SEED

This seed has been treated with INTEGO® SUITE Soybeans which contains clothianidin, ethaboxam, ipconazole and metalaxyl at the rates recommended by the manufacturer.

**DO NOT USE TREATED SEED FOR
FEED, FOOD OR OIL PROCESSING**

**STORE AWAY FROM FEEDS
AND OTHER FOODSTUFFS**

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

- When handling treated seed wear long-sleeved shirt, long pants, socks and shoes, and chemical-resistant gloves: barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinyl chloride or Viton.
- Regardless of application method, do not apply more than 0.2 lb (90.7 g) of clothianidin active ingredient per acre per year. (Based on the seed size of 2800 seeds/lb, at the label rate, the mg of clothianidin will be equal to 0.081 mg/seed).
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.
- Dispose of all excess treated seed. Left over treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements.
- Do not contaminate water bodies when disposing of planting equipment washwaters.
- Dispose of seed packaging in accordance with local requirements.
- Clothianidin is toxic to birds and mammals. Seeds treated with clothianidin exposed on the soil surface may be hazardous to birds and mammals. Cover or collect treated seeds spilled during loading and planting. Dispose of all excess treated seed by burying seed away from bodies of water. Do not contaminate water when disposing of equipment wash-water or rinsate.
- Treated seeds must be planted at a minimum depth of 1 inch.
- This compound is highly toxic to bees exposed directly (contact). Ensure that planting equipment is functioning properly in accordance with manufacturing recommendations to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.



TREATED SEED

This seed has been treated with INTEGO® SUITE Soybeans which contains clothianidin, ethaboxam, ipconazole and metalaxyl at the rates recommended by the manufacturer.

**DO NOT USE TREATED SEED FOR
FEED, FOOD OR OIL PROCESSING**

**STORE AWAY FROM FEEDS
AND OTHER FOODSTUFFS**

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

- When handling treated seed wear long-sleeved shirt, long pants, socks and shoes, and chemical-resistant gloves: barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinyl chloride or Viton.
- Regardless of application method, do not apply more than 0.2 lb (90.7 g) of clothianidin active ingredient per acre per year. (Based on the seed size of 2800 seeds/lb, at the label rate, the mg of clothianidin will be equal to 0.081 mg/seed).
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.
- Dispose of all excess treated seed. Left over treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements.
- Do not contaminate water bodies when disposing of planting equipment washwaters.
- Dispose of seed packaging in accordance with local requirements.
- Clothianidin is toxic to birds and mammals. Seeds treated with clothianidin exposed on the soil surface may be hazardous to birds and mammals. Cover or collect treated seeds spilled during loading and planting. Dispose of all excess treated seed by burying seed away from bodies of water. Do not contaminate water when disposing of equipment wash-water or rinsate.
- Treated seeds must be planted at a minimum depth of 1 inch.
- This compound is highly toxic to bees exposed directly (contact). Ensure that planting equipment is functioning properly in accordance with manufacturing recommendations to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.

Valent U.S.A. LLC
P.O. Box 8025
Walnut Creek CA 94596-8025

IN CASE OF EMERGENCY ONLY: Call 24 Hours A Day 800-892-0099
For **PRODUCT USE** information: Call 800-6-VALENT (800-682-5368)
INTEGO is a registered trademark of Valent U.S.A. LLC

EPA Reg. No. 59639-205
059639-20150717

- This product contains metaxyl which is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.
- Do not graze or feed soybean forage and hay to livestock.
- Areas planted with treated soybean seed may be replanted immediately with bulb vegetables, canola, cereal grains, cotton, fruiting vegetables (including cucurbits), leafy vegetable (including brassica vegetables), rapeseed, root and tuber vegetables and soybean. These areas may also be replanted after 30 days with dry beans, grasses and non-grass animal feeds. These areas may also be replanted after 8 months with sugarcane. Do not plant any other crop in the treated area for at least one year after these treated seeds are planted.
- Do not re-use bags from treated seed to handle food or feed products.
- Do not use augers used for handling treated seed to move seed for feed, food or oil processing.
- Foliar spray of clothianidin can be applied 45 days after planting *INTEGO* SUITE Soybeans treated soybean seed for mid- to late-season foliar insects.
- Not for use on soybeans in California.
- Ipcnazole may persist in soil, and long term use in the same field may result in accumulation of active ingredient.

Valent U.S.A. LLC
P.O. Box 8025
Walnut Creek CA 94596-8025

IN CASE OF EMERGENCY ONLY: Call 24 Hours A Day 800-892-0099
For **PRODUCT USE** information: Call 800-6-VALENT (800-682-5368)
INTEGO is a registered trademark of Valent U.S.A. LLC

EPA Reg. No. 59639-205
059639-20150717

- This product contains metaxyl which is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.
- Do not graze or feed soybean forage and hay to livestock.
- Areas planted with treated soybean seed may be replanted immediately with bulb vegetables, canola, cereal grains, cotton, fruiting vegetables (including cucurbits), leafy vegetable (including brassica vegetables), rapeseed, root and tuber vegetables and soybean. These areas may also be replanted after 30 days with dry beans, grasses and non-grass animal feeds. These areas may also be replanted after 8 months with sugarcane. Do not plant any other crop in the treated area for at least one year after these treated seeds are planted.
- Do not re-use bags from treated seed to handle food or feed products.
- Do not use augers used for handling treated seed to move seed for feed, food or oil processing.
- Foliar spray of clothianidin can be applied 45 days after planting *INTEGO* SUITE Soybeans treated soybean seed for mid- to late-season foliar insects.
- Not for use on soybeans in California.
- Ipcnazole may persist in soil, and long term use in the same field may result in accumulation of active ingredient.