



NipsIt INSIDE[®]

Insecticide

TREATED SEED

This seed has been treated with NipsIt INSIDE[®] Insecticide which contains clothianidin.

**DO NOT USE TREATED SEED
FOR FEED, FOOD OR OIL PROCESSING**

**STORE AWAY FROM FEEDS AND
OTHER FOODSTUFFS**

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

- Wear long-sleeved shirt, long pants and chemical-resistant gloves when handling treated seed.
- Regardless of application method, do not apply more 0.2 lb active ingredient clothianidin/A/year.
- *NipsIt INSIDE* Insecticide is to be used only as a dry-seeded rice seed treatment. Cover the planted seed thoroughly by soil. NipsIt INSIDE Insecticide is not to be followed by a spray treatment of another neonicotinoid insecticide. Follow approved guidelines established for insect resistance management by consulting your local extension pest management expert.
- Do not plant *NipsIt INSIDE* Insecticide treated rice in areas where rice/crawfish aquaculture practices are in place or in or near fish farms, shrimp, prawn or crab pond or nursery operations.
- Rapeseed greens and seed grown or harvested from *NipsIt INSIDE* Insecticide treated seed must not be used for feed or human consumption. Rapeseed grown and harvested from *NipsIt INSIDE* Insecticide treated seed is only for industrial uses and cannot be used for edible oil or any human/feed consumption.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.
- Dispose of all excess treated seed. Left over treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements. Do not contaminate water bodies when disposing of planting equipment washwaters.
- Dispose of seed packaging in accordance with local requirements.
- This compound is highly toxic to bees exposed directly (contact). Ensure that planting equipment is functioning properly in accordance with manufacturing recommendations to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.



NipsIt INSIDE[®]

Insecticide

TREATED SEED

This seed has been treated with NipsIt INSIDE[®] Insecticide which contains clothianidin.

**DO NOT USE TREATED SEED
FOR FEED, FOOD OR OIL PROCESSING**

**STORE AWAY FROM FEEDS AND
OTHER FOODSTUFFS**

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

- Wear long-sleeved shirt, long pants and chemical-resistant gloves when handling treated seed.
- Regardless of application method, do not apply more 0.2 lb active ingredient clothianidin/A/year.
- *NipsIt INSIDE* Insecticide is to be used only as a dry-seeded rice seed treatment. Cover the planted seed thoroughly by soil. NipsIt INSIDE Insecticide is not to be followed by a spray treatment of another neonicotinoid insecticide. Follow approved guidelines established for insect resistance management by consulting your local extension pest management expert.
- Do not plant *NipsIt INSIDE* Insecticide treated rice in areas where rice/crawfish aquaculture practices are in place or in or near fish farms, shrimp, prawn or crab pond or nursery operations.
- Rapeseed greens and seed grown or harvested from *NipsIt INSIDE* Insecticide treated seed must not be used for feed or human consumption. Rapeseed grown and harvested from *NipsIt INSIDE* Insecticide treated seed is only for industrial uses and cannot be used for edible oil or any human/feed consumption.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.
- Dispose of all excess treated seed. Left over treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements. Do not contaminate water bodies when disposing of planting equipment washwaters.
- Dispose of seed packaging in accordance with local requirements.
- This compound is highly toxic to bees exposed directly (contact). Ensure that planting equipment is functioning properly in accordance with manufacturing recommendations to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.

For **PRODUCT USE** information:
 Call 1-800-6-VALENT (1-800-682-5368)
 Call 24 Hours A Day 1-800-892-0099
Valent U.S.A. LLC
 P.O. Box 8025
 Walnut Creek California 94596-8025

EPA Reg. No. 59639-151
Nipsit INSIDE is a registered trademark of Valent U.S.A. LLC



- Do not re-use bags from treated seed to handle food or feed products.
- The compound is toxic to birds and mammals. Treated seeds exposed on soil surface may be hazardous to birds and mammals. Cover or collect treated seeds spilled during loading.
- Pelleted leafy, romaine and head lettuce seeds treated with rates higher than 0.4 mg a/seed must be planted at a minimum depth of 1/2 inch. All other seeds must be planted at a minimum depth of 1/2 inch.
- Do not graze or feed soybean forage and hay to livestock. Treated seed does not flow as freely as untreated seed. Make adjustments to compensate for this restricted seed flow to prevent lower than desired seeding rates.
- **Plant Back Restrictions**
- Areas planted with treated seed of barley, broccoli, buck-wheat, field corn, leafy greens (subgroup 4A), pearl millet, popcorn, proso millet, oats, rice, rye, sugar beet, soybean, teosinte, triticale, and wheat may be replanted immediately with bulb vegetables, canola, cereal grains, cotton, fruiting vegetables (including cucurbits), leafy vegetables (including Brassica vegetables), rapeseed, root and tuber vegetables, sorghum and soybean. These areas may also be replanted after 30 days with dry beans, grasses and non-grass animal feeds. These areas may also be replanted after 8 months with sugarcane. Do not plant any other crop in the treated area for at least one year after these treated seeds are planted.
- Areas planted with treated seed of canola, rapeseed, sorghum and sweet corn may be replanted immediately with bulb vegetables, canola, cereal grains, cotton, fruiting vegetables (including cucurbits), leafy vegetables (including Brassica vegetables), rapeseed, root and tuber vegetables, and soybean. These areas may also be replanted after 30 days with dry beans, grasses and non-grass animal feeds. Do not plant any other crop in the treated area for at least 4 months after these treated seeds are planted.

For **PRODUCT USE** information:
 Call 1-800-6-VALENT (1-800-682-5368)
 Call 24 Hours A Day 1-800-892-0099
Valent U.S.A. LLC
 P.O. Box 8025
 Walnut Creek California 94596-8025

EPA Reg. No. 59639-151
Nipsit INSIDE is a registered trademark of Valent U.S.A. LLC



- Do not re-use bags from treated seed to handle food or feed products.
- The compound is toxic to birds and mammals. Treated seeds exposed on soil surface may be hazardous to birds and mammals. Cover or collect treated seeds spilled during loading.
- Pelleted leafy, romaine and head lettuce seeds treated with rates higher than 0.4 mg a/seed must be planted at a minimum depth of 1/2 inch. All other seeds must be planted at a minimum depth of 1/2 inch.
- Do not graze or feed soybean forage and hay to livestock. Treated seed does not flow as freely as untreated seed. Make adjustments to compensate for this restricted seed flow to prevent lower than desired seeding rates.
- **Plant Back Restrictions**
- Areas planted with treated seed of barley, broccoli, buck-wheat, field corn, leafy greens (subgroup 4A), pearl millet, popcorn, proso millet, oats, rice, rye, sugar beet, soybean, teosinte, triticale, and wheat may be replanted immediately with bulb vegetables, canola, cereal grains, cotton, fruiting vegetables (including cucurbits), leafy vegetables (including Brassica vegetables), rapeseed, root and tuber vegetables, sorghum and soybean. These areas may also be replanted after 30 days with dry beans, grasses and non-grass animal feeds. These areas may also be replanted after 8 months with sugarcane. Do not plant any other crop in the treated area for at least one year after these treated seeds are planted.
- Areas planted with treated seed of canola, rapeseed, sorghum and sweet corn may be replanted immediately with bulb vegetables, canola, cereal grains, cotton, fruiting vegetables (including cucurbits), leafy vegetables (including Brassica vegetables), rapeseed, root and tuber vegetables, and soybean. These areas may also be replanted after 30 days with dry beans, grasses and non-grass animal feeds. Do not plant any other crops in the treated area for at least 4 months after these treated seeds are planted.