





Product Use Overview





| U.S. (excluding California) |

	Product	Key Problems / Uses	Product Benefits and Use Recommendations
Insecticides	Belay® Insecticide	Leafhoppers, mealybugs, <i>Phylloxera</i> spp. (suppression), sharpshooters	Soil Application Long-lasting, broad spectrum insect control that is effective in all soil types and readily available for uptake. Labeled rate range: 6–12 fl oz/A per year. Regardless of the application method, do not apply more than 0.2 lb clothianidin per acre per year. REI: 12 hours. PHI: 30 days.
		Beetles (including Japanese beetle and multi-colored Asian lady beetle), grape berry moth, leafhoppers, mealybugs, sharpshooters	Foliar Application Contact / Ingestion activity with translaminar properties for reliable, broad spectrum control of beetles, leafhoppers, mealybugs and more. Labeled rate range: 2–4 fl oz/A for beetles and leafhoppers; 4–6 fl oz/A for sharpshooters; and 6 fl oz/A for moths and mealybugs. Regardless of the application method, do not apply more than 0.2 lb clothianidin per acre per year. Do not apply this product while bees are foraging. REI: 12 hours. PHI: 0 days.
	Danitol® 2.4 EC Spray	Cutworms, glassy-winged sharpshooters, grape berry moth, Japanese beetle, leafroller, leafhoppers, mites, orange tortrix, plum curculio, rose chafer, spiders, stink bugs (excluding brown)	Quick, broad spectrum pyrethroid insecticide proven not to flare mites. Labeled rate range: 5½–10½ fl oz/A for leafhoppers; 16–21½ fl oz/A for plum curculio; and 10½–21½ fl oz/A for other pests. REI: 24 hours. PHI: 21 days. Danitol is a restricted use pesticide.
	DiPel® DF Biological Insecticide Dry Flowable	Lepidoptera larvae (caterpillar and worm) pests including European grapevine moth (<i>Lobesia botrana</i>), grape berry moth, grape leafroller, grapeleaf skeletonizer (ground only), light brown apple moth, obliquebanded leafroller	<i>DiPel</i> DF is an excellent choice because it delivers effective and economical control of worm pests. And, it has no harvest residue or MRL concerns because it is exempt from tolerances. <i>DiPel</i> DF is a great rotational or tank mix partner to reduce the potential of worms developing resistance to insecticides with other modes of action. Apply 0.5–2 lb/A when larvae are young (early instars) before the crop is damaged. Larvae must be actively feeding on treated, exposed plant surfaces. REI: 4 hours. PHI: 0 days. 
	Esteem® Ant Bait	Ants	For long-lasting control of protein-feeding ants only (such as fire ants). Apply 1.5–2 lb/A at the first signs of protein-feeding ants foraging and actively moving across the vineyard. REI: 12 hours. PHI: 1 day.
	Knack® Insect Growth Regulator	Lecanium scale	Highly-effective IGR for proven, long-lasting scale control. Apply when crawlers first appear. Two applications per season max. Labeled rate: 16 fl oz/A. REI: 12 hours. PHI: 21 days.
	Pyganic® Crop Protection EC 1.4_{II} / 5.0_{II}	Aphids, leafhoppers, stink bugs	Apply when target pest(s) threshold populations are observed. Labeled rate range: <i>Pyganic</i> 1.4 _{II} : 16–64 fl oz/A. <i>Pyganic</i> 5.0 _{II} : 4.5–15.61 fl oz/A. Maximum 10 applications per season. REI: 12 hours. PHI: 0 days. 
	Senstar® Insecticide	Grape tumid gallmaker, lecanium scale (suppression), mealybugs, phylloxera, whiteflies	With two effective modes of action, <i>Senstar</i> provides fast, long-lasting control of soft-bodied insects at all life stages, including eggs. Labeled rate range: 12–16 fl oz/A. REI: 24 hours. PHI: 21 days.
	Venom® Insecticide	Glassy-winged sharpshooters, leafhoppers, mealybugs, <i>Phylloxera</i> (suppression), thrips	Soil Application Super-systemic action with rapid protection against sucking and chewing pests. Labeled rate range: 5–7.5 oz/A. Make only one soil application per season. Do not apply more than a total of 7.5 oz (0.328 lb AI) per acre per season. REI: 12 hours. PHI: 28 days.
		Leafhoppers, mealybugs, multi-colored Asian lady beetle, sharpshooters	Foliar Application Quick control of yield-depleting pests. Labeled rate range: 1–3 oz/A. Do not apply more than a total of 6 oz (0.263 lb AI) per acre per season. REI: 12 hours. PHI: 1 day.
	Zeal® Miticide¹	Mites—European red mite, Pacific, two-spotted, Willamette	Long-lasting mite control with translaminar and ovicidal activity. One application per season. <i>Zeal</i> is a mite growth regulator that affects all mite life stages, including eggs. Labeled rate range: 2–3 oz/A per season. REI: 12 hours. PHI: 14 days.



Product Use Overview



	Product	Key Problems / Uses	Product Benefits and Use Recommendations
Fungicide	Intuity® Fungicide	Botrytis bunch rot, powdery mildew (suppression)	Delivers preventive and residual Botrytis control, locally systemic and translaminar activity, and powdery mildew suppression. Labeled rate: 6 fl oz/A. REI: 12 hours. PHI: 10 days.
Herbicides	Chateau® EZ Herbicide	More than 90 tough broadleaf and grass weeds, including annual morningglory, fleabane, marestail	<i>Chateau EZ</i> is a more convenient, easy-to use liquid formulation. Long-lasting, preemergence herbicide. Won't leach or co-distillate. For table grapes apply between final harvest up budbreak. In juice, raisin and wine grapes you may apply after budbreak only if using a shielded sprayer and spray drift will not contact fruit or foliage. Labeled rate range: 6–12 fl oz/A. Maximum 24 fl oz per year. Clean out sprayer thoroughly after each use. REI: 12 hours. PHI: 60 days.
	Select Max® Herbicide with Inside Technology™ (non-bearing)	Annual and perennial grasses	Superior postemergence for systemic grass control with outstanding crop safety. Lower volatile organic compound (VOC) formulation than other clethodim formulations. May be used with an NIS. Labeled rate range: 9–16 fl oz/A. REI: 24 hours. PHI: 1 year.
Plant Growth Regulators	ProGibb® LV Plus Plant Growth Regulator Solution	Cluster elongation / thinning and increased berry size	Best <i>ProGibb</i> formulation for superior uptake and absorption. Volatile organic compound (VOC) compliant liquid formulation. See product labels for rates and timings by variety. REI: 4 hours. PHI: 0 days. 
	ProTone® SG Plant Growth Regulator Soluble Granule	Enhances red coloration of grapes resulting in earlier harvest, improved harvest management and improved pack-out yield, as well as acceleration of leaf senescence and abscission	Apply 375–1,250 grams/A within the period from 1 week before veraison (50 percent of the target fruit has softened) until anticipated harvest of target fruit. Early timing: First application should be made between 1 week before veraison and 3 weeks after veraison. Late timing: If additional color development is desired, up to three applications can be used. Multiple applications are expected to improve color on cultivars for which there is an extended period between veraison and harvest. Acceleration of leaf senescence and abscission: Apply 469–1,875 grams/A per application. Make one to two foliar applications in combination with other defoliating materials in late fall before the onset of dormancy when defoliation is required. Allow 7–14 days between applications to observe the effect of the first application. REI: 4 hours. PHI: 0 days. <i>ProTone</i> is exempt from a requirement for a tolerance. 



Products That Work, From People Who Care® | valent.com | 800-6-VALENT (682-5368)

Always read and follow label instructions.

with *Inside Technology* is a trademark and *Chateau*, *Esteem*, *Knack*, *Select Max*, *Senstar*, *Venom*, *Zeal* and *Products That Work, From People Who Care* are registered trademarks of Valent U.S.A. LLC. *Select Max* is covered under U.S. Patent No. 7,651,977. *DiPel*, *ProGibb* and *ProTone* are registered trademarks of Valent BioSciences LLC. *Belay*, *Danitol* and *Intuity* are registered trademarks of Sumitomo Chemical Company, Ltd. *Pyganic* is a registered trademark of McLaughlin Gormley King Company. ©2022 Valent U.S.A. LLC. All rights reserved. Printed in the U.S.A. 2022-GRP-8001-R1 10/22

DiPel DF, ProGibb LV Plus, ProTone and Pyganic are OMRI listed and NOP compliant for organic production. 

Danitol is a restricted use pesticide.