

Chateau® EZ Herbicide—A Proven Preemergence Herbicide

Chateau® EZ Herbicide delivers residual preemergence weed control in fallow beds for long-lasting clean fields with rotational flexibility.

- ▶ **Chateau EZ** is a more convenient, **easy-to-use liquid formulation**
- ▶ **Broad Spectrum:** controls more than 90 annual broadleaf and grass weeds, including tough weeds like shepherd's purse, chickweed, lambsquarters, mallow, fleabane and marestalk
- ▶ **Long-Lasting:** stays where you put it; won't leach or co-distillate; no photodegradation
- ▶ **Weather-Ready:** remains active until ¼-inch rainfall or irrigation releases its activity

Crop	Rate (fl oz/A)	Application Timing
Barley	2	3 months before planting
Corn / Fallow	2–4	<ul style="list-style-type: none"> • 30 days before planting at 2–3 fl oz rate • 4 months before planting at 4 fl oz rate
Cotton (no-till / strip-till)	1–2	<ul style="list-style-type: none"> • 14 days before planting at 1 fl oz rate • 21 days before planting at 2 fl oz rate
Cotton (conventional till) / Fallow	2–4	<ul style="list-style-type: none"> • 30 days before planting at 2 fl oz rate • 2 months before planting at 3 fl oz rate • 4 months before planting at 4 fl oz rate
Rice	2	30 days before planting
Rye	2	3 months before planting
Sorghum	2	30 days before planting
Sunflowers	2	30 days before planting
Wheat	2	30 days before planting

How To Use

Rate	Rate determines length of residual weed control and rotation interval
Burn Down	Always tank mix <i>Chateau EZ</i> with the highest rate of a burndown partner, such as glyphosate or other registered herbicide, to control emerged weeds
Timing	Flexible—use to keep fallow beds clean in fall, winter and spring
Method	Ground or air
Spray Volume	10–30 gallons/A—preemergence; 15–60 gallons/A if applying with a burndown herbicide and emerged weeds are present
Adjuvant	Per tank mix partner's recommendation
REI	12 hours

Rotational Restrictions

Rate	Crop	Rotation Interval
1 fl oz/A	Cotton (no-till or strip-till only)	14 days ¹
1.5–2 fl oz/A	Cotton (no-till or strip-till only)	21 days ¹
2 fl oz/A or less	Peanuts, soybeans, sugarcane, sweet potatoes	Immediately
	Field corn (minimum and no-till)	7 days
	Cotton and field corn (conventional tillage), rice, sorghum, sunflowers, tobacco, wheat	30 days ¹
	Barley, dry and snap beans, flax, peas, rye, safflower, sweet corn	3 months
	Alfalfa, canola, clover, oats, potatoes, sugar beets and all other crops not listed ²	<ul style="list-style-type: none"> • 4 months if soil is tilled prior to planting • 8 months if no tillage is performed
	Lentils	6 months
Up to 3 fl oz/A	Peanuts, soybeans, sugarcane and sweet potatoes	Immediately
	Field corn (minimum and no-till)	14 days
	Field corn (conventional tillage), sorghum	30 days ¹
	Cotton, rice, sunflowers, tobacco, wheat	2 months ¹
	Barley, dry and snap beans, flax, pea, rye, safflowers, sweet corn	4 months
	Alfalfa, clover, oats, potatoes, sugar beets	<ul style="list-style-type: none"> • 5 months if soil is tilled prior to planting • 10 months if no tillage is performed
	Canola and all other crops not listed ²	<ul style="list-style-type: none"> • 6 months if soil is tilled prior to planting • 12 months if no tillage is performed
	Lentils	7 months
Up to 4 fl oz/A	Sugarcane	Immediately
	Alfalfa, canola, potatoes, sugar beets and all other crops not listed ²	<ul style="list-style-type: none"> • 6 months if soil is tilled prior to planting • 12 months if no tillage is performed
	Cotton, field corn, peanuts, rice, sorghum, soybeans, sunflowers, tobacco and wheat	4 months
	Transplanted on raised beds only: melons, peppers, tomatoes	2 months (if the top 4 inches of the beds have been removed)
6–12 fl oz/A	Cotton, field corn, peanuts, rice, sorghum, soybeans, sunflowers, tobacco, wheat	9 months
	Alfalfa, canola, sugar beets and all other crops not listed ² Trees can be transplanted 2 months after an application of <i>Chateau EZ</i> ³	<ul style="list-style-type: none"> • 12 months if soil is tilled prior to planting • 18 months if no tillage is performed

¹At least one inch of rainfall/irrigation must occur between application and planting or crop injury may occur.

²Successful soil bioassay must be performed prior to planting these crops.

³Transplanted apples, apricots, avocados, bushberries (including blueberries), cherries, figs, grapes, grapefruit, lemon, nectarine, olives, oranges, peaches, pears, plums (including dried plums), tangerines and tree nuts (including pistachios) can be planted 2 months after a *Chateau EZ* application of 2–12 fl oz/A.