




| Peach Twig Borer Control—California |

## **Outstanding Peach Twig Borer Control**

DiPel® DF Biological Insecticide Dry Flowable is your best choice for control of peach twig borer (PTB) and a broad spectrum of other Lepidopteran pests, especially during flowering when pollinator productivity is critical.

- > Proven, Effective PTB Control: Data shows DiPel DF is highly effective on peach twig borer
- Highly Selective—Won't Harm Beneficial Insects: DiPel DF is compatible in cropping systems and surrounding areas that utilize bee hives for pollination services. DiPel DF is also soft on beneficial insects, workers and the environment.
- DiPel DF is NOP compliant and OMRI Listed for use in organic production
   OMRI

## DiPel DF Peach Twig Borer Control at Bloom



## How To Use

Rate	1 lb/A
Timing	<ul> <li>Apply when larvae are small and actively feeding. For best results, apply at 20–40% egg hatch and again at 80% larval emergence from the hibernacula.</li> <li>Can be utilized during blooming times and does not have any restrictive language for pollinators</li> </ul>
Method	Ground or air
Spray Volume	Coverage is key to control. Use sufficient spray volume to cover all sides of plant surfaces. For aerial applications, apply in 15–20 gallons for large trees and 10–15 gal/A for smaller trees.
Adjuvant	Use a spreader-sticker for hard-to-wet crops or to improve weather-fastness
REI / PHI	4 hours / 0 days (no MRL restrictions)





## Other Important Information

- > Treat when larvae are young (early instars) before the crop is damaged or insects have burrowed into bark
- > Larvae must be actively feeding on treated, exposed plant surfaces to ingest DiPel DF
- Spray foliar to uniformly deposit *DiPel* DF where larvae feed
- If attempting to control a pest with a single spray, treat when egg hatch is essentially complete, but before crop damage occurs



