

Power Pistachio Yields with Quash[®] Fungicide Disease Control

Quash[®] Fungicide is a highly effective foundation fungicide that **delivers proven, broad spectrum performance for a planned rotation program.**

- ▶ Broad spectrum performance
 - Highly effective on Alternaria late blight, Botrytis blossom and shoot blight, Botryosphaeria panicle and shoot blight
- ▶ Preventive protection for when diseases strike
- ▶ Helps deliver on optimal yields and crop quality

Disease Control Ratings



Alternaria Late Blight

Alternaria Late Blight ★★★★★

Botryosphaeria Panicle & Shoot Blight ★★★★★(★)

Botrytis Blossom & Shoot Blight ★★★★★

★★★★★ = excellent and consistent ★★★★★ = good and reliable

Efficacy ratings are from the UC 2022 Efficacy and Treatment Timing of Fungicides for Deciduous Tree Fruit and Nut, Citrus, Strawberry and Vine Crops in California.

How To Use


Rate	4 oz/A per application; maximum 16 oz/A per year
Timing	Apply preventively. Begin treatments when conditions favor disease development. Repeat at 2- to 3-week intervals. Under high disease pressure, use the shorter spray interval.
Method	Foliar spray (100–400 gal/A by ground, 10 gal/A by air)
Rainfast	2 hours
REI / PHI	12 hours / 25 days

Other Important Information

- ▶ Signal word: Caution
- ▶ *Quash* is not hazardous to arthropods, including honeybees
- ▶ FRAC Group 3 Fungicide

Use Timing

Legend: Effectiveness of Treatment Timing



Note: the higher the bar, the more effective *Quash* is at controlling that particular disease at that particular time



Disease	Dormant	April	May	June	July	August	UC-IPM <i>Quash</i> Efficacy Rating
Alternaria Leaf Spot				high	high	medium	★★★★★
Botryosphaeria Panicle & Shoot Blight	low	high	high	high	high	medium	★★★★(★)
Botrytis Blossom & Shoot Blight		high					★★★★