

| Southern |

Regiment® EZ Herbicide Controls Grasses, Broadleaves & Aquatics

Regiment® EZ Herbicide is a contact herbicide that controls yield-robbing grasses, broadleaves and aquatic weeds at a very low use rate. *Regiment EZ* can even control barnyardgrass that is propanil- and/or Facet®-resistant.

- ▶ Controls aquatics, broadleaves and grasses in rice production
- ▶ Great tank mix partner with other herbicides

Control Weeds in Rice Fields with *Regiment EZ*



Untreated



Regiment EZ

How To Use

| | |
|--------------------------|--|
| Timing & Rate | <p>Early postemergence application:</p> <ul style="list-style-type: none"> • 0.33 fl oz/A in combination with a preemergence herbicide at 2-leaf stage • Followed by 1–1.5 fl oz/A either just prior to permanent flood or early post flood <p>Mid postemergence application:</p> <ul style="list-style-type: none"> • 0.9 fl oz/A at 3+ leaf stage • Followed by 0.9 fl oz/A either just prior to permanent flood or early post flood |
| Method | Air or ground |
| Spray Volume | Air: no less than 10 gallons/A Ground: a minimum of 15–20 gallons/A |
| REI | 12 hours |

Other Important Information

- ▶ Include a surfactant, unless otherwise specified on the label. Refer to the Valent bulletin “Approved Surfactants for use with *Regiment* EZ Herbicide” for a list of approved surfactants and rates.
- ▶ Apply to actively growing grasses and weeds
- ▶ DO NOT apply during temperature inversions
- ▶ Medium grain varieties may be more sensitive to *Regiment* EZ than long grain varieties
- ▶ DO NOT apply to the rice variety Bengal
- ▶ May be used on Clearfield[®] and hybrid varieties
- ▶ DO NOT use on the first-year rice on cut ground
- ▶ DO NOT tank mix with malathion, methyl parathion, propanil or propanil-containing mixtures
- ▶ DO NOT make an application of methyl parathion or malathion within 7 days of a *Regiment* EZ application

