

| Fall Application to Control Winter Annuals—Corn, Soybeans, Wheat & More |


Proven Residual Control for your Fall Application

Valor® SX Herbicide offers growers long-lasting residual control of tough weeds like marestail and kochia with minimal rotation restrictions, making it the perfect fit in a fall burndown program. *Valor* SX provides a strong residual herbicide foundation in the fall that:

- ▶ Controls marestail, kochia and other winter annual weeds
- ▶ **Allows** growers to plant earlier in the spring due to cleaner fields
- Decreases loss of moisture and nutrients due to weed competition
- **Lowers** insect pressure from reduced habitat

Valor SX—Residual Control of Kochia in Soybeans





Fall Valor SX followed by spring POST

POST only

Valor SX applied in late October; planted early June the following year.



Valor SX Fall Burndown Efficacy Comparison Data

	Valor SX/ Valor® EZ Herbicide²	Basis® Blend	Autumn™ Super	Metribuzin	Canopy® EX	Simazine
Rotational flexibility	Excellent	Fair	Fair	Excellent	Poor	Poor
Residual marestail control	Excellent	Fair*	Fair*	Not labeled	Fair*	Not labeled
Control of emerged marestail	Good**	Good*	Good*	Good	Fair*	Not labeled
Interactions with other herbicides	No	Yes	Yes	No	No	No
Soil restrictions	Few	Several	Several	Several	Several	Few
Grazing restrictions	0 days#	30 days	45 days	60 days	14 days	Do not graze
Groundwater advisory	No	No	Yes	Yes	No	Yes

^{*}ALS sensitive only—will not control ALS resistant biotypes.

How To Use

Rate & Timing	 2–4 oz/A Application: October–February Best results are obtained when application is made after soil temperature falls below 50°F at a 2-inch depth 			
Tank Mix Partners	Glyphosate, 2,4-D and/or dicamba			
Rotational Flexibility	2 oz/A or less	Field Corn (minimum and no-till): 7 days Field Corn (conventional tillage), Wheat: 30 days ¹ Soybeans: immediately		
	Up to 3 oz/A	Field Corn (minimum and no-till): 14 days Field Corn (conventional tillage): 30 days ¹ Soybeans: immediately Wheat: 2 months ¹		
	Up to 4 oz/A	Field Corn, Soybeans, Wheat: 4 months		

¹At least one inch of rainfall/irrigation must occur between application and planting or crop injury may occur.





[#]When applied to corn stalks after harvest.

^{**}When applied with 2,4-D/dicamba.