

## Chateau® EZ Herbicide—A Proven Preemergence Herbicide

Chateau® EZ Herbicide delivers residual preemergence weed control in fallow beds for long-lasting clean fields with rotational flexibility.

- ▶ **Chateau EZ** is a more convenient, **easy-to-use liquid formulation**
- ▶ **Broad Spectrum:** controls more than 90 annual broadleaf and grass weeds, including tough weeds like shepherd's purse, chickweed, lambsquarters, mallow, fleabane and marestalk
- ▶ **Long-Lasting:** stays where you put it; won't leach or co-distillate; no photodegradation
- ▶ **Weather-Ready:** remains active until ¼-inch rainfall or irrigation releases its activity

Crop	Rate (fl oz/A)	Application Timing
Barley	2	3 months before planting
Corn / Fallow	2–4	<ul style="list-style-type: none"> <li>• 30 days before planting at 2–3 fl oz rate</li> <li>• 4 months before planting at 4 fl oz rate</li> </ul>
Cotton (no-till / strip-till)	1–2	<ul style="list-style-type: none"> <li>• 14 days before planting at 1 fl oz rate</li> <li>• 21 days before planting at 2 fl oz rate</li> </ul>
Cotton (conventional till) / Fallow	2–4	<ul style="list-style-type: none"> <li>• 30 days before planting at 2 fl oz rate</li> <li>• 2 months before planting at 3 fl oz rate</li> <li>• 4 months before planting at 4 fl oz rate</li> </ul>
Rice	2	30 days before planting
Rye	2	3 months before planting
Sorghum	2	30 days before planting
Sunflowers	2	30 days before planting
Wheat	2	30 days before planting

## How To Use

<b>Rate</b>	Rate determines length of residual weed control and rotation interval
<b>Burn Down</b>	Always tank mix <i>Chateau EZ</i> with the highest rate of a burndown partner, such as glyphosate or other registered herbicide, to control emerged weeds
<b>Timing</b>	Flexible—use to keep fallow beds clean in fall, winter and spring
<b>Method</b>	Ground or air
<b>Spray Volume</b>	10–30 gallons/A—preemergence; 15–60 gallons/A if applying with a burndown herbicide and emerged weeds are present
<b>Adjuvant</b>	Per tank mix partner's recommendation
<b>REI</b>	12 hours

## Rotational Restrictions

Rate	Crop	Rotation Interval
1 fl oz/A	Cotton (no-till or strip-till only)	14 days <sup>1</sup>
1.5–2 fl oz/A	Cotton (no-till or strip-till only)	21 days <sup>1</sup>
2 fl oz/A or less	Peanuts, soybeans, sugarcane, sweet potatoes	Immediately
	Field corn (minimum and no-till)	7 days
	Cotton and field corn (conventional tillage), rice, sorghum, sunflowers, tobacco, wheat	30 days <sup>1</sup>
	Barley, dry and snap beans, flax, peas, rye, safflower, sweet corn	3 months
	Alfalfa, canola, clover, oats, potatoes, sugar beets and all other crops not listed <sup>2</sup>	<ul style="list-style-type: none"> <li>• 4 months if soil is tilled prior to planting</li> <li>• 8 months if no tillage is performed</li> </ul>
	Lentils	6 months
Up to 3 fl oz/A	Peanuts, soybeans, sugarcane and sweet potatoes	Immediately
	Field corn (minimum and no-till)	14 days
	Field corn (conventional tillage), sorghum	30 days <sup>1</sup>
	Cotton, rice, sunflowers, tobacco, wheat	2 months <sup>1</sup>
	Barley, dry and snap beans, flax, pea, rye, safflowers, sweet corn	4 months
	Alfalfa, clover, oats, potatoes, sugar beets	<ul style="list-style-type: none"> <li>• 5 months if soil is tilled prior to planting</li> <li>• 10 months if no tillage is performed</li> </ul>
	Canola and all other crops not listed <sup>2</sup>	<ul style="list-style-type: none"> <li>• 6 months if soil is tilled prior to planting</li> <li>• 12 months if no tillage is performed</li> </ul>
	Lentils	7 months
Up to 4 fl oz/A	Sugarcane	Immediately
	Alfalfa, canola, potatoes, sugar beets and all other crops not listed <sup>2</sup>	<ul style="list-style-type: none"> <li>• 6 months if soil is tilled prior to planting</li> <li>• 12 months if no tillage is performed</li> </ul>
	Cotton, field corn, peanuts, rice, sorghum, soybeans, sunflowers, tobacco and wheat	4 months
	Transplanted on raised beds only: melons, peppers, tomatoes	2 months (if the top 4 inches of the beds have been removed)
6–12 fl oz/A	Cotton, field corn, peanuts, rice, sorghum, soybeans, sunflowers, tobacco, wheat	9 months
	Alfalfa, canola, sugar beets and all other crops not listed <sup>2</sup> Trees can be transplanted 2 months after an application of <i>Chateau EZ</i> <sup>3</sup>	<ul style="list-style-type: none"> <li>• 12 months if soil is tilled prior to planting</li> <li>• 18 months if no tillage is performed</li> </ul>

<sup>1</sup>At least one inch of rainfall/irrigation must occur between application and planting or crop injury may occur.

<sup>2</sup>Successful soil bioassay must be performed prior to planting these crops.

<sup>3</sup>Transplanted apples, apricots, avocados, bushberries (including blueberries), cherries, figs, grapes, grapefruit, lemon, nectarine, olives, oranges, peaches, pears, plums (including dried plums), tangerines and tree nuts (including pistachios) can be planted 2 months after a *Chateau EZ* application of 2–12 fl oz/A.