

| Cucurbits, Fruiting, Leafy & Brassica Vegetables—Arizona & California |

Downy Mildew, Pythium and Phytophthora Control

Presidio[®] Fungicide is targeted chemistry that protects crops threatened by downy mildew, damping off and the Phytophthora root, crown and fruit rots caused by *Phytophthora capsici*.




- ▶ **Unique Mode of Action (FRAC Group 43)**—valuable program rotation tool
- ▶ **Effective**—highly systemic movement through the plant
- ▶ **Flexible**—broad spectrum soil-borne and foliar control

Control of Lettuce Downy Mildew

Presidio provided control of downy mildew in lettuce that was equal to the standards

Treatments with the same letter do not differ statistically
Source: Average of six trials



How To Use

Rate	<ul style="list-style-type: none"> • Cucurbits: 4 fl oz/A • Fruiting, leafy and brassica vegetables: 3–4 fl oz/A Consult label for specific application rates
Timing	Consult label for recommended spray timings and intervals
Method	<ul style="list-style-type: none"> • Soil: apply via injection (drip irrigation) • Foliar: apply via ground or aerial application • Adjuvant or Surfactant: none required
Diseases by Crop	<ul style="list-style-type: none"> • Brassica (cole) leafy vegetables: downy mildew, damping off, Phytophthora • Cucurbit vegetables: downy mildew, Phytophthora blight/crown rot • Fruiting vegetables: late blight, Phytophthora root rot, Phytophthora blight • Leafy vegetables (except brassica vegetables): downy mildews, white rust

How To Use continued

Resistance Management	<ul style="list-style-type: none"> • Tank mix <i>Presidio</i> with another fungicide with a different mode of action against the same disease • For cucurbit downy mildew, mancozeb 3 lb/A or chlorothalonil 2 pt/A is recommended • After <i>Presidio</i> + tank mix application, rotate to a non-Group 43 fungicide 	
Restrictions	Maximum applications <ul style="list-style-type: none"> • Cucurbits: 2 • Fruiting, leafy and brassica vegetables: 4 	Sequential applications <ul style="list-style-type: none"> • Cucurbits: None • Fruiting, leafy and brassica vegetables: 2
PHI	2 days in cucurbits, fruiting, leafy and brassica vegetables	
REI	12 hours	

Rotational Restrictions

The following rotational crops may be planted after applying *Presidio* at the labeled rate. Planting earlier than the labeled rotational interval is not allowed.

Crops		Rotational Interval
Basil	Leafy Vegetables except Brassica	Immediately
Brassica (Cole) Leafy Vegetables (Crop Group 5)	Vegetables (Crop Group 4)	
Bulb Vegetables (Crop Group 3-07)	Listed Succulent Beans	
Citrus Fruit (Crop Group 10-10)	Root and Tuber Vegetables (Crop Group 1)	
Cucurbit Vegetables (Crop Group 9)	Small Fruits, Vine Climbing, except	
Fruiting Vegetables (Crop Group 8-10)	Fuzzy Kiwifruit (Crop Subgroup 13-07F)	
Hops	Tobacco	
Field Corn	Wheat	30 days
All Other Crops		18 months

Other Important Information

- ▶ Group 43 fungicide: sole fungicide in FRAC Group 43
- ▶ Formulation: 4 SC
- ▶ Packaging: 10 quarts/case
- ▶ *Presidio* is compatible with many fungicides and insecticides
- ▶ Rainfast after 2 hours
- ▶ PPE: long sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material, socks and shoes