



METLOCKTM

FUNGICIDE

TREATED SEED

This seed has been treated with METLOCKTM Fungicide containing metconazole.

CAUTION
**DO NOT USE FOR FEED,
FOOD OR OIL PROCESSING**
**STORE AWAY FROM FEEDS AND
OTHER FOODSTUFFS**

KEEP OUT OF REACH OF CHILDREN

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear long-sleeved shirt, long pants and waterproof gloves when handling treated seed.

FIRST AID

If swallowed: Call a poison control center or doctor immediately for treatment advice.
Have a person sip a glass of water if able to swallow.
Do not induce vomiting unless told to do so by a poison control center or doctor.
Do not give anything to an unconscious person.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes.
Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
Call a poison control center or doctor for treatment advice.

- Dispose of all excess treated seed. Left over treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements. Do not contaminate water bodies when disposing of planting equipment washwaters.
- Dispose of seed packaging in accordance with local requirements.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading.
- No plant back restriction for Barley, Canola, Corn (including field, sweet and pop) Oat, Peanut, Rye, Soybean, Sugar Beet, Triticale and Wheat. A 30-day plant back interval for brassica leafy vegetables and leafy vegetables. A 120-day plant back interval for all other crops.

EPA Reg. No. 59639-171

METLOCK is a trademark of Valent U.S.A. Corporation

For **PRODUCT USE** information:
Call 1-800-6-VALENT (1-800-682-5368)
IN CASE OF EMERGENCY ONLY
Call 24 Hours A Day 1-800-892-0099

Valent U.S.A. Corporation
P.O. Box 8025
Walnut Creek, California 94596-8025