

| Soybeans\* |

## Improve Consistency of Postemergence Herbicides with Resource® Herbicide

Resource® Herbicide improves control of velvetleaf, lambsquarters, waterhemp and other pigweeds as compared to glufosinate, dicamba, 2,4-D and other postemergence herbicides applied alone in a wide variety of soybean trait platforms.

- ▶ Postemergence herbicide partner including approved tank mix partner with XtendiMax® and Enlist One® herbicides
- ▶ Improves control of problematic weeds and increases speed of activity for postemergence program
- ▶ Can be applied regardless of your soybean trait platform (excluding organic)
- ▶ Included in the John Deere Financial program

### How To Use

<b>Rate</b>	<p><b>Spring Burndown:</b> 2–4 fl oz/A, + tank mix partner</p> <p><b>Postemergence:</b> 4–12 fl oz/A + tank mix partner</p> <ul style="list-style-type: none"> <li>• Do not apply more than 12 fl oz /A in a single application or more than a total of 16 fl oz /A to soybeans during a single growing season</li> <li>• Do not apply if rain is expected within one hour of application; otherwise unsatisfactory weed control may result</li> </ul>
<b>Timing</b>	Burndown or postemergence up to 60 days prior to harvest (follow restrictions associated with tank mix partner)
<b>Spray Volume</b>	15–20 gal/A
<b>Adjuvant</b>	<ul style="list-style-type: none"> <li>• NIS 0.25% v/v or COC 1 pt/A</li> <li>• Refer to tank mix partner label</li> <li>• Valent recommends use of a Chemical Producers &amp; Distributors Association-certified adjuvant</li> </ul>
<b>Rotation Restrictions</b>	Do not rotate to crops other than field corn or soybeans within 30 days after last application
<b>Hay / Grazing Restrictions</b>	Do not graze treated fields or harvest for forage or hay
<b>REI / PHI</b>	12 hours / 60 days

**\*Not for use in California on soybeans.**