

Power Pistachio Yields with Quash[®] Fungicide Disease Control

Quash[®] is a highly effective foundation fungicide that **delivers proven, broad spectrum performance for a planned rotation program.**

- ▶ Broad spectrum performance
 - Highly effective on Alternaria late blight, Botrytis blossom and shoot blight, Botryosphaeria panicle and shoot blight
- ▶ Preventive protection for when diseases strike
- ▶ Helps deliver on optimal yields and crop quality

Disease Control Ratings



Alternaria Late Blight

Alternaria Late Blight ★★★★★

Botryosphaeria Panicle & Shoot Blight ★★★

Botrytis Blossom & Shoot Blight ★★★(★)

★★★★ = excellent and consistent ★★★ = good and reliable

Efficacy ratings are from the UC 2017 Efficacy and Treatment Timing of Fungicides for Deciduous Tree Fruit, Nut, Strawberry and Vine Crops publication

How To Use


Rate	4 oz/A per application; maximum 16 oz/A per year
Timing	Apply preventively. Begin treatments when conditions favor disease development. Repeat at 2- to 3-week intervals. Under high disease pressure, use the shorter spray interval.
Method	Foliar spray (100–400 gal/A by ground, 10 gal/A by air)
Rainfast	2 hours
PHI / REI	25 days / 12 hours

Other Important Information

- ▶ Signal word: Caution
- ▶ *Quash* is not hazardous to arthropods, including honeybees
- ▶ FRAC Group 3 Fungicide

Use Timing

Legend: Effectiveness of Treatment Timing



Note: the higher the bar, the more effective *Quash* is at controlling that particular disease at that particular time



Disease	Dormant	April	May	June	July	August	UC-IPM <i>Quash</i> Efficacy Rating
Alternaria Leaf Spot				High	High	Medium	★★★★
Botryosphaeria Panicle & Shoot Blight	Low	High	High	High	High	Medium	★★★
Botrytis Blossom & Shoot Blight		High					★★★★(★)