



Senstar™ Insecticide: Boost Your Insect Control

New Senstar™ Insecticide brings two effective modes of action for fast, long-lasting control of soft-bodied insects including whiteflies, aphids, psyllids and thrips. With translaminar and systemic activity that affects all life stages of targeted pests for long-lasting control, *Senstar* helps ensure high-quality produce from the start.

- ▶ Controls target pests at all life stages, including eggs
- ▶ Provides dual systemic activity to control pests moving into new vegetative growth as well as pests which populate in already developed foliage
- ▶ Delivers fast translaminar movement to reach target pests that feed on the underside of leaves
- ▶ Provides high selectivity to harmful insect species with no hazardous effects on many beneficial insects for excellent fit in Integrated Pest Management (IPM) programs

Senstar Delivers Powerful Whitefly Control at All Life Stages



Senstar provided faster and more consistent control of eggs, nymphs and adults than *Knack* or *Movento*.

* Number of individuals per leaf (8 days after second application)

Means followed by the same letters are not significantly different ($P \leq 0.10$)

Source: St. Lucie County, FL; Better Crops, LLC

Senstar delivered faster control of eggs and nymphs of whiteflies seven days after initiating the insecticidal program to control whiteflies.

** Eggs and nymphs (5 leaves/plant); adults (10 leaves/plant evaluated)

Means followed by the same letters are not significantly different ($P \leq 0.10$)

Source: St. Lucie County, FL; Better Crops, LLC

Senstar Vegetable Crops and Pest Summary

| Crops | Pests | Rate (fl oz/A) | PHI (days) |
|---|---|-------------------|--|
| Brassica (Cole) Leafy Vegetables | Aphids, diamondback moth*, swedge midge, thrips (larvae)*, whiteflies | 6–10 | 7 |
| Bulb Vegetables | Onion thrips (larvae), Western flower thrips* | 10 | 3 (subgroup 3-07A) 7 (subgroup 3-07b) |
| Carrots | Aphids, whiteflies | 10 | 7 |
| Fruiting Vegetables | Aphids, leafminers*, psyllids, Western flower thrips (larvae)*, whiteflies | 8–10 | 1 |
| Leafy Vegetables | Aphids, diamondback moth*, leafminers*, Western flower thrips (larvae)*, whiteflies | 8–10 | 14 |
| Sugar Beets | Aphids, root maggot*, whiteflies | 9–18 | 28 |
| Tuberous & Corm Vegetables | Aphids, psyllids, Western flower thrips (larvae)* | 8–10 | 7 |
| Watercress | Aphids, sharpshooters, whiteflies | 8–24 | 7 |

*Suppression only

Other Important Information

- ▶ Do not apply more than 10 fl oz/A per application (18 fl oz/A per application for sugar beets; 24 fl oz/A per application for watercress)
- ▶ Do not make more than 2 applications per crop season
- ▶ Do not apply more than 20 fl oz/A per crop season (36 fl oz/A per crop season for sugar beets; 48 fl oz/A per crop season for watercress)
- ▶ Minimum interval between applications is 14 days

Integrated Pest Management (IPM)

Senstar is highly suited for use in IPM programs because it shows high selectivity to harmful insect species with no hazardous effects on many beneficial insects.