


Downy Mildew, Pythium and Phytophthora Control

Presidio[®] Fungicide is targeted chemistry that protects crops threatened by downy mildew, damping off and the Phytophthora root, crown and fruit rots caused by *Phytophthora capsici*.

- ▶ **Unique Mode of Action (FRAC Group 43)**—valuable program rotation tool
- ▶ **Effective**—highly systemic movement through the plant
- ▶ **Flexible**—broad spectrum soil-borne and foliar control



Control of Lettuce Downy Mildew



Presidio provided control of downy mildew in lettuce that was equal to the standards

Treatments with the same letter do not differ statistically

Source: Average of six trails

Other Important Information

- ▶ Group 43 fungicide: sole fungicide in FRAC Group 43
- ▶ Formulation: 4 SC
- ▶ Packaging: 10 quarts/case
- ▶ Presidio is compatible with many fungicides and insecticides
- ▶ Rainfast after 2 hours
- ▶ PPE: long sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material, socks and shoes

How To Use

Rate	<ul style="list-style-type: none"> • Cucurbits: 4 fl oz/A • Fruiting, leafy and brassica vegetables: 3–4 fl oz/A Consult label for specific application rates		
Timing	Consult label for recommended spray timings and intervals		
Method	<table border="1"> <tr> <td> <ul style="list-style-type: none"> • Soil: apply via injection (drip irrigation) • Foliar: apply via ground or aerial application • Adjuvant or Surfactant: none required </td> <td> Water volume: <ul style="list-style-type: none"> • Ground application: 20–100 gal/A • Aerial application: 5 gal/A minimum </td> </tr> </table>	<ul style="list-style-type: none"> • Soil: apply via injection (drip irrigation) • Foliar: apply via ground or aerial application • Adjuvant or Surfactant: none required 	Water volume: <ul style="list-style-type: none"> • Ground application: 20–100 gal/A • Aerial application: 5 gal/A minimum
<ul style="list-style-type: none"> • Soil: apply via injection (drip irrigation) • Foliar: apply via ground or aerial application • Adjuvant or Surfactant: none required 	Water volume: <ul style="list-style-type: none"> • Ground application: 20–100 gal/A • Aerial application: 5 gal/A minimum 		
Resistance Management	<ul style="list-style-type: none"> • Tank mix <i>Presidio</i> with another fungicide with a different mode of action against the same disease • For cucurbit downy mildew, mancozeb 3 lb/A or chlorothalonil 2 pt/A is recommended • After <i>Presidio</i> + tank mix application, rotate to a non-Group 43 fungicide 		
Restrictions	Maximum applications <ul style="list-style-type: none"> • Cucurbits: 2 • Fruiting, leafy and brassica vegetables: 4 Sequential applications <ul style="list-style-type: none"> • Cucurbits: None • Fruiting, leafy and brassica vegetables: 2 		
PHI	2 days in cucurbits, fruiting, leafy and brassica vegetables		
REI	12 hours		