



| Rice–Southern |

Early Control of Tough Broadleaf, Aquatic and Sedge Weeds

League® Herbicide is an unprecedented early season, lead-off play against tough broadleaf, aquatic and sedge weeds. It offers solid preemergence residual control of yield-robbing weeds and will fit your herbicide program lineup.

- ▶ Strikes weeds as they emerge providing unique residual control and also offers postemergence control for best fit in your weed program
- ▶ **Controls tough rice weeds**, including jointvetches, hemp sesbania, Texasweed, morningglory, flat sedge, eclipta, yellow nutsedge and more
- ▶ Fit for use on conventional or Clearfield® rice

How To Use

Preemergence	<ul style="list-style-type: none"> • Rate: <i>League</i> 6.4 oz/A plus foundation grass herbicide (such as <i>Bolero</i>® Herbicide, <i>Command</i>® or <i>Newpath</i>®) • Residual broadleaf and sedge control—yellow nutsedge, early indigo (jointvetch), coffeebean and Texasweed control • Good approach for broadleaf control around towns and soybeans • Not for use on clay soils with pH >8.0
Postemergence (Preflood) (2- to 3-leaf rice)	<ul style="list-style-type: none"> • Option 1: Early post—<i>League</i> 4 oz/A plus <i>Bolero</i>, <i>Command</i>, <i>Facet</i>®, <i>Prowl</i>® or <i>Newpath</i> • Option 2: Cleanup prior to flood—<i>League</i> 4 oz/A plus <i>Regiment</i>® Herbicide, <i>Newpath</i>, <i>Facet</i> or propanil (weed spectrum dependent) • Contact (<2-inch weeds) and residual • Weeds—Pennsylvania smartweed, coffeebean, indigo, ducksalad, yellow nutsedge, Texasweed • For grass control (especially barnyardgrass) mix with <i>Regiment</i>, <i>Facet</i> or propanil. Do not mix with <i>Ricestar</i>® HT or <i>Clincher</i>®.
Sequential (Preemergence)	<i>League</i> 3.2 oz/A plus foundation grass herbicide (such as <i>Bolero</i> , <i>Command</i> or <i>Newpath</i>)
Early Postemergence (Preflood)	<ul style="list-style-type: none"> • <i>League</i> 3.2 oz/A plus a postemergence product such as <i>Regiment</i>, <i>Newpath</i>, <i>Facet</i> or propanil • For grass control (especially barnyardgrass), mix with <i>Regiment</i>, <i>Facet</i> or propanil. Do not mix with <i>Ricestar</i> HT or <i>Clincher</i>.
Method	Ground or air
Spray volume	15–20 gal/A (ground); 10 gal/A (air)
REI	12 hours



Weeds Controlled by League

Broadleaves & Aquatics		Grasses Tank mix with a foundation grass herbicide, such as Bolero, Command, Newpath	
Cutleaf groundcherry	<i>Physalis angulata</i>	Barnyardgrass	<i>Echinochloa crus-galli</i>
Ducksalad	<i>Heteranthera</i> spp.	Broadleaf signalgrass	<i>Brachiaria platyphylla</i>
Eclipta	<i>Eclipta prostrata</i>	Crabgrass, large	<i>Digitaria sanguinalis</i>
Gourd, Texas	<i>Cucurbita foetidissima</i>	Johnsongrass (seedling)	<i>Sorghum halepense</i>
Hemp Sesbania	<i>Sesbania exaltata</i>	Panicum, Fall	<i>Panicum dichotomiflorum</i>
Jointvetch, Indian	<i>Aeschynomene indica</i>	Panicum, Texas	<i>Panicum texanum</i>
Jointvetch, Northern	<i>Aeschynomene virginica</i>	Sprangletop	<i>Leptochloa</i> spp.
Morningglory, pitted	<i>Ipomoea lacunosa</i>	Sedges	
Pigweed	<i>Amaranthus</i> spp.	Dayflower	<i>Commelina communis</i>
Redstem	<i>Ammannia</i> spp.	Flatsedge, rice	<i>Cyperus iria</i>
Texasweed	<i>Caperonia palustris</i>	Nutsedge, yellow	<i>Cyperus esculentus</i>
Woolly Croton	<i>Croton capitatus</i>		

Other Important Information

- ▶ *League* is approved for aerial application. Do not apply *League* by air within 0.5 miles of emerged cotton or non-STS soybeans.
- ▶ In Arkansas for aerial application, *League* can be tank mixed with Command as follows: *League* + Command + Newpath or RiceBeaux® or SuperWHAM!® or Clearpath® or Grasp® SC
- ▶ *League* can be tank mixed with any rice herbicides except Ricestar HT or Clincher
- ▶ Do not apply to second crop (stubble/ratoon) rice
- ▶ Fields with a history of weed resistance to Londax® may exhibit resistance to *League*
- ▶ Do not apply to stressed rice
- ▶ Do not drain the field for 7 days after application when making a postemergence application to a flooded field
- ▶ Do not apply more than 6.4 oz of *League* per acre per year
- ▶ For tank mix applications with propanil-containing products, read and follow the entire label of each product to be used in the tank mix
- ▶ Do not apply an organophosphate insecticide within 21 days before, or 7 days after, an application of *League*



Products That Work, From People Who Care® | valent.com | 800-6-VALENT (682-5368)

Always read and follow label instructions.

League and *Products That Work, From People Who Care* are registered trademarks of Valent U.S.A. LLC. *Bolero* and *Regiment* are registered trademarks of Kumiai Chemical Industry Co., Ltd. *Clincher* and *Grasp* are registered trademarks of The Dow Chemical Company. *Command* is a registered trademark of FMC Corporation. *Clearfield*, *Clearpath*, *Facet*, *Newpath* and *Prowl* are registered trademarks of BASF. *Londax* is a registered trademark of United Phosphorus, Inc. *RiceBeaux* and *SuperWHAM!* are registered trademarks of RiceCo LLC. *Ricestar* is a registered trademark of Bayer. ©2018 Valent U.S.A. LLC. All rights reserved. Printed in the U.S.A. 2018-LEA-8000 LF 12/18