

Chateau® Herbicide SW—A Proven Preemergence Residual Herbicide

- ▶ **Broad Spectrum:** controls **more than 90** annual broadleaf and grass weeds, including marestail, fleabane, mallow, filaree, annual ryegrass and other annual grasses
- ▶ **Long-Lasting:** stays where you put it; won't leach or co-distillate; no photodegradation
- ▶ **Weather-Ready:** remains active until ¼-inch rainfall or irrigation releases its activity



Photo shows excellent weed control in spring from fall application of *Chateau* at 12 oz/A

How To Use

Rate	<ul style="list-style-type: none"> • 6–12 oz/sprayed acre. For maximum residual control, use the higher rate. Apply with burndown herbicide if emerged weeds are present. • If weed pressure is heavy, make two sequential applications. Wait at least 30 days between applications. • Maximum 24 oz/A per year
Method	Spray a uniform band directed at the base of the trunk.
Spray Volume	10–30 GPA
Timing	Apply after final harvest and before bloom in spring for greatest crop safety. Apply to apple before silver tip. Apply to stone fruit and pear before bud break.
Tank Mix Partners	Cheetah®, glyphosate, Gramoxone Inteon®, Treevix®, Rely® (glufosinate), etc., add burndown power to control emerged weeds. <i>Chateau</i> can also be tank mixed with other preemergence herbicides such as Prowl®, Alion® and Karmex® to extend weed spectrum as needed following most restrictive labeling.
REI / PHI	12 hours / 60 days



Get a Head Start on Weed Control with *Chateau*

	<i>Chateau</i>	Matrix®	Princep®	Alion	Karmex
Apple (Tree Age)	Anytime*	>1 year	>1 year	>3 years	>1 year
Cherry (Tree Age)	Anytime*	>1 year	>3 years	>3 years	Not registered
Peach (Tree Age)	Anytime*	>1 year	>3 years	>3 years	>3 years

*With protection from spray contact if <1 year old

Other Important Information—See Label for Complete Information

- ▶ *Chateau* requires at least ¼ inch of rainfall or irrigation for activation
- ▶ For young trees (<3 years) in sandy soils (sand + gravel >80%), apply up to 6 oz/A per season
- ▶ Apply carefully—although *Chateau* is primarily a preemergence herbicide, direct spray contact or drift will damage foliage, green bark and fruit
 - Do not apply to trees established <1 year unless protected from spray with non-porous wrap, grow tubes or waxed containers
- ▶ Do not apply within 300 yards of non-dormant pears
- ▶ Do not apply to farm alleys or roads where traffic may result in treated dust settling onto crops or other desirable vegetation
- ▶ Use a spray drift agent with all *Chateau* applications. Add AMS to spray tank before mixing *Chateau* and a spray drift agent. Conduct a jar test before mixing.
- ▶ Clean out spray tank daily
- ▶ **Oregon and Washington—Specific Information**
 - Do not apply to pear orchards
 - Use **only** in the following counties:
 - Oregon:** Benton, Clackamas, Clatsop, Columbia, Coos, Curry, Douglas, Jackson, Josephine, Lane, Lincoln, Linn, Marion, Morrow, Multnomah, Polk, Tillamook, Umatilla, Yamhill and Washington
 - Washington:** Benton, Clallam, Clark, Cowlitz, Franklin, Grant, Grays Harbor, King, Jefferson, Kitsap, Lewis, Pacific, Pierce, Skagit, Snohomish, Thurston, Wahkiakum, Walla Walla and Whatcom